

The Jan Mayen Microplate Complex and the Wilson Cycle

Christian Schiffer^{1*}, Alexander Peace², Jordan Phethean¹, Laurent Gernigon³, Ken McCaffrey¹, Kenni D. Petersen⁴, Gillian Foulger¹

¹*Department of Earth Sciences, Durham University, DH1 3LE, UK*

²*Department of Earth Sciences, Memorial University of Newfoundland, St. John's, NL, A1C 5S7, Canada*

³*Geological Survey of Norway (NGU), Trondheim, Leiv Erikssons vei 39, Norway*

⁴*Department of Geoscience, Aarhus University, 8000 Aarhus, Denmark*

**Corresponding Author (email: christian.schiffer@zoho.com)*

Abstract

The opening of the North Atlantic region was one of the most important geodynamic events that shaped the present-day passive margins of Europe, Greenland and North America. Although well-studied, much remains to be understood about the evolution of the North Atlantic, including the role of the Jan Mayen Microplate Complex (JMMC). Geophysical data provide an image of the crustal structure of this microplate and enable a detailed reconstruction of the rifting and spreading history. However, the mechanisms that cause separation of microplates between conjugate margins are still poorly understood. In this contribution, we assemble recent models of rifting and passive margin formation in the North Atlantic and discuss possible scenarios that may have led to formation of the JMMC. This event has likely been triggered by regional plate-tectonic reorganisations rejuvenating inherited structures. The axis of rifting and continental breakup and the width of the JMMC was controlled by old Caledonian fossil subduction/suture zones. Its length is related to E-W oriented deformation and fracture zones possibly linked to rheological heterogeneities inherited from pre-existing Precambrian terrane boundaries.

(end of abstract)

The North Atlantic region inspired some aspects of plate tectonic theory (Fig. 1). These include the Wilson Cycle which predicts the closure of oceans leading to continent-continent collision followed by their reopening along former sutures (Wilson 1966, Dewey & Spall 1975). The North Atlantic is often considered to be a text-book example

33 of an ocean that opened along the former sutures of at least two temporarily distinct
34 orogenic events – the Neoproterozoic Grenvillian-Sveconorwegian and the early
35 Palaeozoic Caledonian-Variscan orogenies (Ryan & Dewey, 1997; Vauchez *et al.*,
36 1997; Bowling & Harry, 2001; Thomas, 2006; Misra, 2016). Nevertheless, some
37 aspects of the North Atlantic geology remain enigmatic, such as the formation of the
38 North Atlantic Igneous Province (NAIP) (Vink, 1984; White & McKenzie, 1989;
39 Foulger & Anderson, 2005; Meyer *et al.*, 2007), the development of the volcanic
40 passive margins (Franke, 2013; Geoffroy *et al.*, 2015), the formation of Iceland and the
41 development of the Jan Mayen Microplate Complex (JMMC), also referred to as the Jan
42 Mayen Microcontinent (Foulger *et al.*, 2003; Gaina *et al.*, 2009; Gernigon *et al.*, 2015).
43 The JMMC comprises both oceanic and continental crust, probably highly thinned and
44 magmatically modified (Kuvaas & Kodaira, 1997; Blischke *et al.*, 2016 and references
45 therein). Large parts of it remain to be studied, however. Other continental fragments
46 have been identified in the North Atlantic region (Nemčok *et al.*, 2016) and more may
47 underlie parts of Iceland and/or the Iceland-Faroe Ridge (Fedorova *et al.*, 2005;
48 Foulger, 2006; Paquette *et al.*, 2006; Gernigon *et al.*, 2012; Torsvik *et al.*, 2015).

49

50 *Geological Setting of the North Atlantic region*

51 Following the collision of Laurentia, Baltica and Avalonia in the Ordovician and
52 Silurian (Roberts 2003, Gee *et al.* 2008, Leslie *et al.* 2008), and subsequent
53 gravitational extensional collapse in the late orogenic phases (Dewey, 1988; Dunlap &
54 Fossen, 1998; Rey *et al.*, 2001; Fossen, 2010), the North Atlantic region experienced
55 lithospheric delamination and associated uplift over a period of 30-40 Ma, followed by
56 a long period of rifting (Andersen *et al.*, 1991; Dewey *et al.*, 1993). Phases of extension
57 and cooling transitioned into continental rifting that led to final continental breakup and
58 seafloor spreading between Greenland and Europe in the early Palaeogene (Talwani &
59 Eldholm 1977, Skogseid *et al.* 2000). During the late Mesozoic, continental breakup
60 propagated simultaneously southward from the Eurasia Basin and northward from the
61 Central Atlantic initially into the Labrador Sea- Baffin Bay rift system and then into the
62 North Atlantic (Srivastava, 1978; Doré *et al.*, 2008). Whether rifting, continental
63 breakup, and associated magmatism was initiated by mantle upwelling, for example a
64 deep mantle plume (White & McKenzie, 1989; Hill, 1991; Nielsen *et al.*, 2002; Rickers
65 *et al.*, 2013) or plate-driven processes (Nielsen *et al.*, 2007; Ellis & Stoker, 2014)

66 (“bottom-up” or “top down” views) is still under debate (van Wijk *et al.*, 2001; Foulger
67 *et al.*, 2005b; Lundin & Doré, 2005; Simon *et al.*, 2009).

68 The North Atlantic spreading axis initially comprised the Reykjanes Ridge, the Aegir
69 Ridge, east of the JMMC and the Mohns Ridge farther north (Talwani & Eldholm,
70 1977; Nunns, 1982, Fig. 1). Independent rotation of the JMMC resulted in fan-shaped
71 opening of the Norway Basin, during the Eocene (Nunns, 1982; Gaina *et al.*, 2009;
72 Gernigon *et al.*, 2012). This reconfiguration led to a second phase of breakup and the
73 separation of the JMMC from Greenland at approximately magnetic anomaly chron C7
74 (~24 Ma) (Vogt *et al.*, 1970; Gaina *et al.*, 2009; Gernigon *et al.*, 2015). After a period of
75 simultaneous rifting on both the Aegir Ridge and the complex JMMC/proto-Kolbeinsey
76 rift/ridge system (Doré *et al.*, 2008; Gaina *et al.*, 2009; Gernigon *et al.*, 2015), the Aegir
77 Ridge was abandoned in the Oligocene and the spreading centre relocated to the west of
78 the JMMC onto the Kolbeinsey Ridge. The present-day North Atlantic shows evidence
79 for a dynamic contribution of the topography, requiring an anomalous pressure anomaly
80 uplifting the lithosphere that might be linked to Iceland (Schiffer & Nielsen, 2016).

81 Although the history of rifting in the North Atlantic is becoming increasingly better
82 constrained, the mechanisms controlling the location, timing, and formation of rifts,
83 fracture zones, and associated microcontinents are still poorly understood. The
84 formation of the JMMC has been traditionally attributed to mantle plume impingement
85 and subsequent lithospheric weakening (Müller *et al.* 2001). More recently it has been
86 suggested to result from the breaching of lithosphere weakened as a result of pre-
87 existing structures (*e.g.*, Schiffer *et al.* 2015b). The final separation of the JMMC is also
88 spatially and temporally linked to enhanced magmatic activity and the subsequent
89 formation of Iceland (Doré *et al.*, 2008; Tegner *et al.*, 2008; Larsen *et al.*, 2013; Schiffer
90 *et al.*, 2015b) but it lacks the classic features of a volcanic passive margin (*e.g.*,
91 underplating, seaward dipping reflectors) along its western continent-ocean boundary,
92 conjugate to the East Greenland margin (Kodaira *et al.*, 1998; Breivik *et al.*, 2012;
93 Peron-Pinvidic *et al.*, 2012; Blischke *et al.*, 2016). In this paper, we discuss the possible
94 role of pre-existing structure and inheritance in formation of the JMMC as an extension
95 to the Wilson Cycle and plate tectonic theory.

96

97 **JAN MAYEN MICROPLATE COMPLEX**

98 The JMMC has a bathymetric signature stretching over 500 km from north to south in
99 the central part of the Norwegian-Greenland Sea (Fig. 1) (Gudlaugsson *et al.* 1988,
100 Kuvaas & Kodaira 1997, Blischke *et al.* 2016). It is bordered to the north by the Jan
101 Mayen Fracture Zone (JMFZ) and the volcanic complex of Jan Mayen Island. To the
102 south, it is bordered by the NE coastal shelf of Iceland which is part of the Greenland-
103 Iceland-Faroe Ridge (GIFR), a zone of shallow bathymetry approximately 1100 km
104 length (Figs. 1 and 2). The JMMC separates the Norway Basin to the east from the
105 Iceland Plateau to the west (Vogt *et al.* 1981, Kandilarov *et al.* 2012, Blischke *et al.*
106 2016).

107 The JMMC crust has been inferred to be continental primarily on the basis of seismic
108 refraction data (Kodaira *et al.*, 1997; Kodaira *et al.*, 1998; Mjelde *et al.*, 2007a; Breivik
109 *et al.*, 2012; Kandilarov *et al.*, 2012). However, for large areas of the JMMC crustal
110 affinity remains uncertain, particularly near Iceland in the south (Breivik *et al.*, 2012;
111 Brandsdóttir *et al.*, 2015) due to the lack of geophysical data and boreholes (see
112 Gernigon *et al.*, 2015 and Blischke *et al.*, 2016 for data coverage). Fundamentally, the
113 distribution of oceanic versus continental crust, as well as the nature of the deformation
114 expected between the JMMC, Iceland and the Faroe continental block are unknown.
115 Recent high-resolution aeromagnetic data and pre-rift reconstructions of the Norwegian-
116 Greenland Sea show that the southern JMMC underwent extreme thinning during the
117 first phase of breakup and, as it now has a width of ~500 km, 400% of extension has
118 occurred compared to its pre-drift configuration (Gernigon *et al.* 2015). It seems
119 unlikely that this extreme extension is entirely accommodated by the thinning of
120 continental crust. We cannot rule out the possibility that the southern JMMC partly
121 comprises igneous crust (Gernigon *et al.*, 2015) or exhumed mantle (Blischke *et al.*,
122 2016).

123 An oceanic fracture zone might be present south of the JMMC between the northeastern
124 tip of the Iceland Plateau and the Faroe Islands in the southeast (i.e. the postulated
125 Iceland-Faroe Fracture Zone, IFFZ, see Fig. 1 and 2, e.g. Blischke *et al.* 2016).
126 However, an oceanic fracture zone or transform requires oceanic lithosphere on both
127 sides and, given the uncertain crustal affinity this interpretation is speculative. A
128 lineament exists north of the Iceland-Faroe Ridge (IFR, the part of the GIFR east of and
129 including Iceland) but magnetic and gravity potential-field data do not provide
130 conclusive evidence for a real oceanic transform or fracture zone (Fig. 3). Gernigon *et*
131 *al.* (2012) showed that continuation of the magnetic chrons mapped in the Norway

132 Basin and the high-magnetic trends observed along the IFR remain unclear, notably due
133 to the low quality, the sparse distribution of the magnetic profiles along the IFR and
134 later igneous overprint related to the formation of Iceland. No magnetic chrons are
135 identified in the broad NE-SW magnetic lineations, especially west of the Faroe
136 Platform. Additional magnetic disparities are associated with lateral variations of
137 basement depth and possible discrete ridge jumps (e.g. Smallwood & White, 2002;
138 Hjartarson et al., 2017). The GIFR comprises anomalous thick crust (>20-25 km)
139 possibly associated with massive crustal underplating, which is generally attributed to
140 increased magmatism (Staples et al., 1997; Richardson et al., 1998; Smallwood et al.,
141 1999; Darbyshire et al., 2000; Greenhalgh & Kusznir, 2007). The origin and nature of
142 the GIFR remains controversial (McBride et al., 2004), also because the crust shows
143 atypical geophysical properties and differs from “normal” continental and oceanic crust
144 (Bott, 1974; Foulger et al., 2003). A recent paper (Hjartarson et al., 2017) favours an
145 oceanic origin of the IFR, but the authors do not exclude the presence of seaward
146 dipping reflectors and old basement in the expected "oceanic domain". Some authors
147 suggested that the excess thickness under Iceland may be partly attributed to buried
148 continental crust possibly extending up to the JMMC and Iceland (Fedorova et al.,
149 2005; Foulger, 2006). Continental zircons and geochemical analysis of lavas in
150 southeast Iceland support the presence of continental material (Paquette et al., 2006;
151 Torsvik et al., 2015). The Aegir Ridge and the Reykjanes Ridge might have never
152 connected during the early stage of spreading of the Norway Basin involving complex
153 overlapping spreading segments along the IFR. Such overlapping spreading ridges may
154 have preserved continental lithosphere in between (Gaina et al., 2009; Gernigon et al.,
155 2012, 2015; Ellis & Stoker, 2014). Ellis & Stoker (2014) suggested that no complete
156 continental breakup along the IFR happened before the separation of the JMMC and the
157 appearance of Iceland (first dated eruptions at ~18 Ma). Gernigon et al. (2015)
158 suggested earlier breakup possibly between C22/C21 (~47 Ma) and C6 (~24Ma) during
159 the onset of significant rifting in the southern part of the JMMC. The continental
160 lithosphere east of Iceland (the IFR, Fig. 1) probably didn't entirely breach in the early
161 rifting of the North Atlantic (e.g. C24r-C22, Early Eocene). To avoid further ambiguity,
162 we refer to it as the Iceland-Faroe accommodation zone (IFAZ). Consequently, the
163 IFAZ may characterize local continental transform margin segments, a diffuse strike-
164 slip fault zone and/or a more complex oblique/transensional continental rift system that
165 initially formed along the trend of the proto IFR.

166 MICROPLATE FORMATION

167 An aspect of the Wilson Cycle that requires more clarification (Thomas, 2006; Huerta &
168 Harry, 2012; Buitter & Torsvik, 2014) is whether the locations of major, pre-existing
169 structures can explain the formation, location and structure of microplates such as the
170 JMMC (Schiffer *et al.* 2015a). Understanding the formation of continental fragments is
171 crucial to understanding continental breakup (Lavie & Manatschal, 2006; Peron-
172 Pinvidic & Manatschal, 2010). Microcontinents and continental ribbons represent one
173 category of continental fragments produced during rifting and breakup (Lister *et al.*,
174 1986; Peron-Pinvidic & Manatschal, 2010; Tetreault & Buitter, 2014).

175 We follow the original definition of a microcontinent Scrutton (1976) that it must
176 contain: (i) pre-rift basement rocks, (ii) crust and lithosphere of continental affinity,
177 horizontally displaced from the original continent and surrounded by oceanic crust, and
178 (iii) a distinct morphological feature in the surrounding oceanic basins. Such a system
179 between two pairs of conjugate margins may also include isolated fragments of oceanic
180 crust and lithosphere that deformed together before final and definitive isolation from
181 the conjugate continents. To make a distinction, we call such a feature a microplate
182 complex, and it can involve several sub-plates of oceanic and/or continental affinity. A
183 true microcontinent will, therefore, comprise just one kind of microplate complex. The
184 most important aspect of the present study is that such a microplate complex, like a true
185 microcontinent, is separated from the main continental conjugate margins by two or
186 more spreading ridges. The cause, history and processes leading to relocalisation of the
187 complex are not well understood. Suggested mechanisms include the impact of a mantle
188 plume (Müller *et al.*, 2001; Gaina *et al.*, 2003; Mittelstaedt *et al.*, 2008), global plate-
189 tectonic reorganisation (Collier *et al.*, 2008; Gaina *et al.*, 2009), and ridge "jumps" that
190 exploit inhomogeneities, weaknesses and rheological contrasts in the continental
191 lithosphere after the abandonment of a previous spreading ridge (Abera *et al.* 2016,
192 Sinha *et al.* 2016). This could be nascent or inherited underplating (Yamasaki &
193 Gernigon 2010) and/or fossil suture zones. Strike-slip mechanisms under different
194 transtensional and transpressional stress regimes have also been proposed to generate
195 microcontinents (Nemčok *et al.* 2016). Microplates can also result from crustal
196 fragmentation during volcanic margin formation by large-scale continent-vergent faults
197 formed/activated by strengthening of the deep continental crust – the so-called "C-
198 Block" mechanism (Geoffroy *et al.* 2015).

199 Whittaker *et al.* (2016) proposed a model for microcontinent formation between
200 Australia and Greater India whereby changes in plate motion direction caused
201 transpression and stress buildup across large-offset fracture zones, leading to transfer of
202 deformation to a less resistive locus (Fig. 4). Their proposed model is as follows.
203 Initially NW-SE spreading separated Australia from Greater India with transtensional or
204 strike-slip motion along the Wallaby-Zenith Fracture Zone from 133 Ma. A plume
205 (Kerguelen) is postulated to have been in the vicinity and may have maintained and/or
206 enhanced crustal weakening of the SE Greater India rifted margin. Reorganisations of
207 motion between Australia and Greater India to a NNW-SSE direction at 105 Ma
208 resulted in transpression along the NW-SE-oriented Wallaby-Zenith Fracture Zone. As
209 a result, the spreading centre relocated to the west along the continental margin of India,
210 calving off the Batavia and Gulden Draak microcontinents, and resulting in
211 abandonment of the Dirck Hartog spreading ridge to the south (Fig. 4).

212

213 **NORTH ATLANTIC – STRUCTURE AND INHERITANCE**

214 The classic Wilson Cycle model envisages closure and reopening of oceans along
215 continental sutures. In this model, breakup is thus guided by lithospheric inheritance
216 from previous orogenesis (Wilson 1966, Dewey & Spall 1975). Inheritance,
217 rejuvenation and control of pre-existing structure on localising deformation occurs on
218 various scales and styles beyond large-scale breakup of continents (Holdsworth *et al.*,
219 1997; Manatschal *et al.*, 2015). Inherited features may include crustal or lithospheric
220 thickness variations, structural and compositional heterogeneity across terrane
221 boundaries, accreted terranes, sedimentary basins and/or intruded, metamorphosed and
222 metasomatised material and fabrics. These heterogeneities may also cause thermal and
223 rheological anomalies that vary in size, depth and degree of anisotropy, that can
224 potentially be rejuvenated given the appropriate stresses (Krabbendam & Barr, 2000;
225 Tommasi *et al.*, 2009; Manatschal *et al.*, 2015; Tommasi & Vauchez, 2015). Inheritance
226 is an important control on rifting, passive-margin end-member style (*e.g.*, volcanic or
227 non-volcanic) (Vauchez *et al.*, 1997; Bowling & Harry, 2001; Chenin *et al.*, 2015;
228 Manatschal *et al.*, 2015; Schiffer *et al.*, 2015b; Svartman Dias *et al.*, 2015; Duretz *et al.*,
229 2016; Petersen & Schiffer, 2016), the formation of fracture zones, transform faults,
230 transform margins (Thomas, 2006; Gerya, 2012; Doré *et al.*, 2015), magmatism
231 (Hansen *et al.* 2009, Whalen *et al.* 2015), compressional deformation (Sutherland *et al.*

232 2000, Gorczyk & Vogt 2015, Heron *et al.* 2016), the breakup of supercontinents and
233 supercontinent cycles (Vauchez *et al.*, 1997; Audet & Bürgmann, 2011; Frizon de
234 Lamotte *et al.*, 2015).

235

236 *Precambrian orogenies*

237 In Canada, Greenland and Northwest Europe, multiple suturing events have built
238 continental lithosphere that comprises Archean-to-early Proterozoic cratons surrounded
239 by younger terranes. Preserved sutures and subduction zones in the interior of the
240 cratons have survived subsequent amalgamation demonstrating that crustal and upper
241 mantle heterogeneities may persist for billions of years (Balling 2000, van der Velden &
242 Cook 2005). Terrane boundaries of any age may act as rheological boundaries that
243 influence or control crustal deformation long after their formation and independently of
244 subsequent plate motions. Major Precambrian terrane boundaries in the North Atlantic
245 region are shown in Figure 2.

246 Multiple Precambrian suturing events have contributed to the amalgamation of the
247 Baltic Shield in Scandinavia. The Lapland-Kola mobile belt formed by accretion of
248 various Archean to Palaeoproterozoic terranes, including the oldest Karelian terrane
249 (Gorbatshev & Bogdanova 1993, Bergh *et al.* 2012, Balling 2013). This was followed
250 by the late Palaeoproterozoic Svecofennian accretion, the formation of the
251 Transscandinavian Igneous Belt, and finally the Meso-Neoproterozoic Sveconorwegian
252 orogeny (Gorbatshev & Bogdanova, 1993; Bingen *et al.*, 2008; Bergh *et al.*, 2012;
253 Balling, 2013; Slagstad *et al.*, 2017).

254 Precambrian terranes are also preserved in Greenland, the oldest of which are Archean
255 in age and include the North Atlantic and Rae Cratons (St-Onge *et al.* 2009). The
256 components that together constitute the North Atlantic Craton formed 3850 – 2550 Ma
257 (Polat *et al.* 2014) and the Rae Craton formed 2730 – 2900 Ma (St. Onge *et al.* 2009).
258 Paleoproterozoic terranes in Greenland surround the North Atlantic Craton and include
259 (i) the Nagssugtoqidian Orogen (Van Gool *et al.* 2002), (ii) the Rinkian Orogen
260 (Grocott & McCaffrey 2016) and (iii) the Ketilidian Mobile Belt (Garde *et al.* 2002).

261 The Precambrian terranes of northeast Canada, Greenland and Scandinavia are thought
262 to have formed as coherent mobile belts (Kerr *et al.*, 1996; Wardle *et al.*, 2002; St-Onge
263 *et al.*, 2009). As Greenland and North America have not undergone significant relative

264 lateral motions or rotation the interpretation of conjugate margins is relatively simple
265 (Kerr *et al.*, 1996; Peace *et al.*, 2016). In contrast, whether or not Baltica has
266 experienced rotation (Gorbatshev & Bogdanova 1993, Bergh *et al.* 2012) is currently
267 unresolved.

268

269 *Caledonian Orogeny*

270 Formation of the Ordovician to Devonian Caledonian-Appalachian Orogen preceded
271 rifting, ocean spreading and subsequent passive margin formation of the present-day
272 North Atlantic. This Himalaya-style orogen involved at least two phases of subduction:
273 (i) the early eastward-dipping Grampian-Taconian event and (ii) the late westward-
274 dipping Scandian event that led to the assembly of part of Pangaea (Roberts 2003, Gee
275 *et al.* 2008). During orogenesis the structural fabric of the crust and lithospheric mantle
276 can be reoriented resulting in fabric anisotropy that localises subsequent deformation
277 (Tommasi *et al.*, 2009; Tommasi & Vauchez, 2015).

278 High-velocity, lower-crustal bodies (HVLCB) are observed along many passive
279 continental margins (Lundin & Doré, 2011; Funck *et al.*, 2016a) and have been
280 traditionally associated with magmatic underplating or intrusions into the lower crust of
281 passive margins during breakup (Olafsson *et al.* 1992, Eldholm & Grue 1994, R. Mjelde
282 *et al.* 2007, White *et al.* 2008, Thybo & Artemieva 2013). However, with improved data
283 alternative interpretations have been proposed such as syn-rift serpentinitisation of the
284 uppermost mantle under passive margins (Ren *et al.*, 1998; Reynisson *et al.*, 2010;
285 Lundin & Doré, 2011; Peron-Pinvidic *et al.*, 2013). It has also been suggested that part
286 of the continental HVLCB may be remnants of inherited metamorphosed crust or
287 hydrated meta-peridotite that existed prior to initial rifting and continental breakup
288 (Gernigon *et al.*, 2004; Gernigon *et al.*, 2006; Fichler *et al.*, 2011; Wangen *et al.*, 2011;
289 Mjelde *et al.*, 2013; Nirrengarten *et al.*, 2014).

290 Mjelde *et al.* (2013) have identified a number of such “orogenic” HVLCB along
291 different parts of the North Atlantic passive margins (the South- and Mid-Norwegian
292 margin, East Greenland margin, SW Barents Sea margin, Labrador margin), which may
293 have higher than normal upper mantle velocities ($V_p > 8.2$ km/s). These may comprise
294 eclogitised crust and be part of the Iapetus Suture. Petersen & Schiffer (2016) proposed
295 a mechanism to explain the presence of old inherited HVLCB beneath the rifted

296 margins and concluded that they could represent preserved and subsequently deformed
297 pre-existing subduction/suture zones that were activated during rifting and continental
298 breakup. Eclogite in a fossil slab has a similar but weaker rheology than the surrounding
299 “dry olivine” lithosphere (after Zhang & Green, 2007), while a fossil, hydrated mantle
300 wedge acts as an effective and dominant weak zone. Eclogites of the Bergen Arcs
301 (Norway) show softening due to fluid infiltration Jolivet *et al.* (2005). These ultra-high
302 velocity HVLCB (ultra-HVLCB) are distributed primarily along the mid-Norwegian
303 margin and the Scoresbysund area in East Greenland (Mjelde *et al.*, 2013). This
304 suggests that at least one fossil subduction zone may have been subject to rift-related
305 deformation and exhumation (Petersen & Schiffer 2016).

306 Structures in the Central Fjord area of East Greenland (Schiffer *et al.* 2014), the Flannan
307 reflector in northern Scotland (Snyder & Flack 1990, Warner *et al.* 1996) and the
308 Danish North Sea (Abramovitz & Thybo 2000) have been interpreted as preserved
309 orogenic structures of Caledonian age (i.e. fossil subduction or suture zones) (Fig. 2).
310 Schiffer *et al.* (2015a) proposed that the Central Fjord structure and the Flannan
311 reflector once formed a contiguous eastward-dipping subduction zone, possibly of
312 Caledonian age, that may have influenced rift, magmatic, and passive-margin evolution
313 in the North Atlantic (Figure 2). Combined geophysical-petrological modelling of the
314 Central Fjord structure suggests it comprises a relict hydrated mantle wedge associated
315 with a fossil subduction zone (Schiffer *et al.* 2015b, Schiffer *et al.* 2016). The most
316 recent Caledonian subduction event was associated with the Scandian phase leading to
317 the westward subduction of Iapetus crust (Roberts 2003, Gee *et al.* 2008). Evidence of
318 this subduction zone in the form of a preserved slab has not been detected in the
319 lithospheric mantle of the Norwegian Caledonides. However, structures in the crust and
320 upper mantle in the Danish North Sea detected by the Mona Lisa experiments
321 (Abramovitz & Thybo 2000) might be the trace of this subduction. HVLC indicative of
322 eclogite along the Mid-Norwegian margin (Mjelde *et al.*, 2013) and Norwegian North
323 Sea (Christiansson *et al.*, 2000; Fichler *et al.*, 2011) might also represent deformed
324 remnants of the Scandian subduction.

325 *Fracture and accommodation zones*

326 The JMMC is bound by two tectonic boundaries including the East and West Jan
327 Mayen Fracture Zones in the north and the postulated Iceland-Faroe accommodation
328 zone (IFAZ) in the south. These tectonic boundaries accommodated and allowed the

329 non-rigid microplate to move independently from the surrounding North Atlantic
330 oceanic domains (Gaina *et al.*, 2009; Gernigon *et al.*, 2012, 2015).

331 Relationships between pre-existing structures and the formation of large-scale shear and
332 fracture zones, oceanic transforms or other accommodation/deformation zones have
333 been proposed in previous work (Mohriak & Rosendahl, 2003; Thomas, 2006; Taylor *et al.*,
334 *et al.*, 2009; de Castro *et al.*, 2012; Gerya, 2012; Bellahsen *et al.*, 2013; Gibson *et al.*,
335 2013). The location, orientation and nature of fracture zones in the North Atlantic may
336 be linked to lithospheric inheritance (Behn & Lin, 2000). For example, the Charlie-
337 Gibbs Fracture Zone between Newfoundland and the British/Irish shelf has been linked
338 to the location of the Iapetus suture and inheritance of compositional and structural
339 weaknesses (Tate 1992, Buitter & Torsvik 2014). The Bight Fracture Zone might be
340 linked to the Grenvillian front, which is exposed in Labrador (Lorenz *et al.* 2012).

341 The IFAZ could represent a complex discontinuity zone along the present-day IFR.
342 Along this transition zone between the Reykjanes, Aegir and Kolbeinsey ridges
343 fragments of continental crust may be preserved together with discontinuous and/or
344 overlapping oceanic fragments later affected by significant magmatic overprint (the
345 Icelandic “swell”, Bott, 1988). In the geodynamic context, it may have formed along the
346 fossil subduction zone proposed to have existed between the East Greenland and
347 British/Irish margins (Fig. 2). It has also been proposed that it may have comprised part
348 of the “Kangerlussuak Fjord tectonic lineament”, a NW-SE-oriented lineament in east
349 Greenland (Tegner *et al.* 2008).

350 Other deformation zones may correlate with Precambrian basement terrane boundaries
351 in Scandinavia. These are overprinted by Caledonian deformation, obscuring older
352 relationships (cf. CDF in Fig. 2) and generating new orogenic fabrics (Vauchez *et al.*,
353 1998). The westward extrapolation of the northern Sveconorwegian suture may
354 correlate with the East Jan Mayen Fracture Zone (EJMFZ), whilst extrapolation of the
355 Svecofennian-Karelian suture may correspond to the formation of the Senja Fracture
356 Zone (SFZ) (Doré *et al.* 1999, Fichler *et al.* 1999, Indrevær *et al.* 2013). Extrapolation
357 of the Karelian-Lapland Kola terrane suture converges with the complex DeGeer
358 Fracture Zone that marks the transition of the North Atlantic to the Arctic Ocean (Engen
359 *et al.* 2008). These correlations suggest that Precambrian basement inheritance localises
360 strain during initial continental rifting. However, the exact location and grade of
361 deformation of Precambrian sutures under the Caledonides and the highly stretched

362 continental margins is often poorly known or not known at all. Thus, any correlation is
363 speculative and requires future work.

364 *Iceland and magmatic evolution*

365 Factors including the thermal state of the crust and mantle, small scale convection,
366 upwelling, composition, volatile content, and lithospheric and crustal structure may all
367 play roles (King & Anderson, 1998; Asimow & Langmuir, 2003; Korenaga, 2004;
368 Foulger *et al.*, 2005a; Hansen *et al.*, 2009; Brown & Leshner, 2014; Chenin *et al.*, 2015;
369 Hole & Millett, 2016).

370 Inheritance may influence the amount of volcanism produced in the North Atlantic
371 because volcanic passive margins preferentially develop in regions of heterogeneous
372 crust where Palaeozoic orogenic belts separate Precambrian terranes. Inversely, magma-
373 poor margins often develop in the interiors of orogenic belts with either uniform-
374 Precambrian or younger-Palaeozoic crust (Bowling & Harry, 2001). For example, the
375 intersection of the East Greenland-Flannan fossil subduction zone with the North
376 Atlantic rift axis correlates spatially and temporally with pre-breakup magmatism, the
377 formation of JMMC and the occurrence of the Iceland melt anomaly along the sub-
378 parallel GIR (Schiffer *et al.*, 2015b).

379 Prior to breakup (ca. 55 Ma), magma was dominantly emplaced along and south-west of
380 the proposed East Greenland-Flannan fossil subduction zone (Fig. 2) (Ziegler, 1990;
381 Torsvik *et al.*, 2002). This may be partly an effect of the south-to-north “unzipping” of
382 the pre-North Atlantic lithosphere. Other processes that produce enhanced mantle
383 melting are increased temperature, mantle composition and active asthenospheric
384 upwelling (Brown & Leshner, 2014). The zonation of areas with and without magmatism
385 may suggest that the proposed structure is a boundary zone between lithospheric blocks
386 of different composition and rheology that react differently to applied stresses. Different
387 relative strength in crust and mantle lithosphere, for instance, could cause depth
388 dependent deformation, where thinning is focussed in the mantle lithosphere (Huisman
389 & Beaumont 2011). Petersen & Schiffer (2016) demonstrated that extension of orogenic
390 lithosphere with thickened crust (>45 km) leads to depth-dependent thinning where the
391 mantle lithosphere breaks earlier than the crust and as a result encourages pre-breakup
392 magmatism. Indirectly, sub-continental mantle heterogeneities may encourage
393 localisation of deformation leading to rapid and sudden increase in lithospheric thinning
394 (Yamasaki & Gernigon, 2010). These processes could contribute to pre-breakup

395 adiabatic decompression melting (Petersen & Schiffer 2016). Enhanced magmatism
396 could also be caused by a lowered solidus due to presence of eclogite (Foulger *et al.*,
397 2005a), water in the mantle (Asimow & Langmuir 2003) or CO₂ (Dasgupta &
398 Hirschmann, 2006). Atypical magmatism is, surprisingly, observed along the
399 interpolated axis of the proposed fossil subduction zone than elsewhere. It currently
400 coincides with the GIFR where igneous crustal thickness is inferred to be greatest (Bott,
401 1983; Smallwood *et al.*, 1999; Holbrook *et al.*, 2001; Mjelde & Faleide, 2009; Funck *et*
402 *al.*, 2016b). However, it is unclear whether the entire thickness of “Iceland type crust”
403 (Bott, 1974; Foulger *et al.*, 2003) has crustal petrology (Foulger *et al.*, 2003; Foulger &
404 Anderson, 2005).

405 Higher water contents have been recorded in basalts and volcanic glass in the vicinity of
406 the fossil subduction zone (the Blossville Kyst, East Greenland, Iceland and one
407 sample from the Faroe Islands, see Fig. 2) than in regions further away from Iceland
408 (West Greenland, Hold with Hope, Reykjanes Ridge) (Jamtveit *et al.* 2001, Nichols *et*
409 *al.* 2002). This is consistent with a hydrated upper mantle source as a consequence of
410 melting Caledonian subducted materials (Schiffer *et al.* 2015a). Water in the mantle
411 may also contribute to enhanced melt production and thus unusually thick igneous crust
412 (Asimow & Langmuir 2003).

413 The formation of the Iceland Plateau (>18 Ma) followed extinction of the Aegir Ridge
414 and full spreading being taken up on the Kolbeinsey Ridge (Dore *et al.* 2008). This
415 spreading ridge migration was contemporaneous with far-field plate tectonic
416 reconfigurations, cessation of seafloor spreading in the Labrador-Baffin Bay system
417 (Chalmers & Pulvertaft 2001) and a global change of Greenland plate motion from SW-
418 NE to W-E (Gaina *et al.*, 2009; Abdelmalak *et al.*, 2012).

419

420 **AN INHERITANCE MODEL FOR FORMATION OF THE JMMC**

421 We propose a new tectonic model for formation of the JMMC that links rejuvenation of
422 old and pre-existing orogenic structures to global plate tectonic reconfigurations. In our
423 model a change in the orientation of the regional stress field in the Eocene rejuvenated
424 pre-existing structures with favourable orientations. This caused localisation of
425 extension and spreading ridges resulting in the formation of a microplate between the
426 large European and American/Greenland continental plates. Our model closely follows

427 that of Whittaker *et al.* (2016), with the extension that a fossil subduction zone is
428 utilised as a physical and compositional weak zone that helps to accommodate a second
429 axis of breakup (Fig. 5). Plate tectonic reorganisations and rejuvenation of pre-existing
430 structures may not be the only controls on continental breakup, but they may be the
431 dominant ones in the case of the JMMC. In areas where no microplate formation is
432 observed continental breakup followed the youngest, weakest Caledonian collision
433 zone, the Scandian, west-dipping subduction in Scandinavia. This may have been better
434 aligned with the ambient stress field during rifting and/or breakup. Following the model
435 of Petersen & Schiffer (2016), the remnants of this subduction zone or other inherited
436 orogenic structures may now be distributed along the Mid-Norwegian margin as pre-
437 breakup HVLCB (Christiansson *et al.*, 2000; Gernigon *et al.*, 2006; Fichler *et al.*, 2011;
438 Wangen *et al.*, 2011; Mjelde *et al.*, 2013; Nirrengarten *et al.*, 2014; Mjelde *et al.*, 2016).
439 The subduction zone was already deformed in the Norwegian North Sea by rifting
440 subsequent to the Permo-Triassic and is still preserved as a large HVLCB beneath the
441 North Sea rift (Christiansson *et al.* 2000, Fichler *et al.* 2011). A stronger, east-dipping
442 subduction zone in East Greenland, may also have been deformed but did not
443 accommodate breakup. Continental rifting and possible overlapping of the Reykjanes
444 and Mohns ridge leading initiating the JMMC formation (Gernigon *et al.*, 2012, 2015)
445 may have been promoted by the presence of this deep-rooted weak zone.

446 The Caledonian and Grenvillian orogenic fabric and major associated structures are
447 generally parallel to the NNE-SSE trend of rifting in the North Atlantic with some
448 exceptions, such as the opening of Labrador Sea. Older terrane boundaries are close to
449 perpendicular. Young Caledonian structures define the axis of rifting and continental
450 breakup. This can be explained by the presence of deep, weak eclogite-facies roots
451 along the axis of the Caledonian Orogen, and extensional collapse of the Caledonian
452 mountain range causing earlier extension to initiate perpendicular to the axis of collision
453 (Ryan & Dewey, 1997; Rey *et al.*, 2001). Precambrian structures are still preserved in
454 stable cratons surrounded by orogens and mobile belts. Once rifting occurs, lateral
455 weaknesses and rheological boundaries control segmentation of the rift axis and
456 eventually influence the formation of across-strike deformation zones of different kinds,
457 *e.g.*, fracture and transform zones, diffuse/oblique/transtensional rift and ridge systems.

458 Our suggested scenario for the formation of the JMMC complements the established
459 Wilson Cycle concept. We propose that reactivation and petrological variation of
460 inherited structures of different ages, coupled with changes in the regional/global stress

461 regime, controlled microplate formation in the following sequence of events (see also
462 Fig. 6):

- 463 1. Early Palaeocene: Rifting propagates from the Central Atlantic into the Labrador
464 Sea - Baffin Bay rift system (Roest & Srivastava, 1989; Chalmers & Pulvertaft,
465 2001; Peace *et al.*, 2016)
- 466 2. Early Eocene (Fig. 6b): Change in Labrador Sea-Baffin Bay spreading direction
467 from NW-SE to W-E (Abdelmalak *et al.*, 2012) and onset of seafloor spreading
468 in the northeast Atlantic (Gaina *et al.*, 2009). This was possibly related to the
469 far-field stress field applied by the collision of Africa and Europe (Nielsen *et al.*,
470 2007) and/or to the relocation of the postulated Iceland plume (Skogseid *et al.*,
471 2000; Nielsen *et al.*, 2002).
- 472 3. The NW-SE stress field in the North Atlantic between Greenland and
473 Scandinavia would have favoured deformation on deep structures associated
474 with the Iapetus Suture on the Norwegian margin rather than the East Greenland
475 margin with the proposed fossil subduction zone (Fig. 2). Thus, initial breakup is
476 generally parallel to and in the vicinity of the Iapetus Suture.
- 477 4. The Iceland-Faroe Accommodation Zone (IFAZ) forms as the southern limit of
478 the JMMC and may be linked to localisation of strain along the proposed fossil
479 subduction zone or other potential rheological boundaries. No continental
480 breakup occurred between Iceland and the Faroe Islands (Iceland Faroe Ridge),
481 with underlying, uninterrupted but thinned, continental lithosphere (Ellis &
482 Stoker, 2014).
- 483 5. Mid-late Eocene: Accelerated extension occurred in the southern part of the
484 JMMC and local reorganisation of the Norway Basin spreading system
485 (Gernigon *et al.* 2012, 2015) developed around 47 Ma (Fig. 6c) A first phase of
486 magmatism between Greenland and the proto-JMMC was initiated (Tegner *et*
487 *al.*, 2008; Larsen *et al.*, 2014). In the southern JMMC, isolated spreading cells
488 possibly developed before steady state development of the Kolbeinsey Ridge.
- 489 6. Late Eocene - early Oligocene (Fig. 6c): A major plate tectonic reorganisation
490 including a change from NW-SE to NE-SW plate motion coincident with
491 abandonment of seafloor spreading along the Labrador Sea-Baffin Bay system
492 and consequent cessation of anti-clockwise rotation of Greenland (Mosar *et al.*,
493 2002; Gaina *et al.*, 2009; Oakey & Chalmers, 2012). This change in plate motion
494 results in deformation along the fracture zones and transpression on the IFAZ.

- 495 7. Locking of the IFAZ triggered continental breakup between Greenland and the
496 proto-JMMC subsequent to continental rifting between them. This is consistent
497 with the microplate model of Whittaker *et al.* (2016) for the Indian Ocean.
498 Rotational rifting between Greenland and the proto-JMMC started much earlier
499 (c. 47-48 Ma) than abandonment of the Labrador Sea-Baffin Bay spreading
500 system (c. 40 Ma) and breakup between Greenland and the JMMC (33-24 Ma).
- 501 8. Ultraslow spreading continued on the Aegir Ridge after ca. 31 Ma (Mosar *et al.*,
502 2002; Gaina *et al.*, 2009; Gernigon *et al.*, 2015), while drastic rifting and
503 possible embryonic spreading developed south of the proto-JMMC until steady
504 state spreading along Kolbeinsey Ridge was completely established at 24 Ma
505 (Vogt *et al.*, 1970; Doré *et al.*, 2008; Gernigon *et al.*, 2012).
- 506 9. The Aegir Ridge was abandoned with all plate motion accommodated by the
507 Kolbeinsey Ridge after 24 Ma, separating the proto-JMMC from East Greenland
508 (Fig 6d). The West Jan Mayen Fracture Zone, the eastern branch of which had
509 already been established during the opening of the Norway Basin, then
510 connected the Kolbeinsey Ridge with the Mohns Ridge north of the JMMC.

511 **SUMMARY**

512

513 We propose a new model for formation of a microplate complex as an extension to the
514 established Wilson Cycle concept. The new model invokes rejuvenation of major pre-
515 existing structures by plate-driven processes controlling both breakup and JMMC
516 formation.

517

518 The initial axis of continental breakup exploited lithospheric weaknesses associated
519 with the Iapetus Suture (Fig. 6 a,b). These structures were particularly susceptible to
520 deformation due to their preferential orientation with respect to the NW-SE to W-E
521 oriented extensional stress field. Fracture zones and strike-slip/oblique zones of
522 deformation delineate the later-forming JMMC. The IFAZ represents one of these zones
523 and may have formed along an old subduction zone. The origin of the IFAZ remains
524 poorly defined because of poor data coverage. However, it is likely that despite extreme
525 thinning of the continental lithosphere no continental breakup occurred between
526 present-day JMMC and the Faroe Islands (e.g. Gernigon *et al.*, 2015; Blischke *et al.*,
527 2016).

528

529 Our model predicts that, following a major change in extension direction that was
530 coeval with the abandonment of the Labrador Sea-Baffin Bay oceanic spreading and
531 transform system, oblique deformation occurred south of the proto-JMMC and along
532 the poorly defined IFAZ (Fig. 6c). This caused further westward relocation of the
533 spreading centre towards a fossil subduction zone where eclogite and, especially, weak
534 inherited serpentinite accommodated the relocation and final development of the
535 Kolbeinsey Ridge. Complete development of the Kolbeinsey Ridge resulted in final
536 separation of the proto-JMMC from East Greenland (Fig. 6d) and complete breakup of
537 the North Atlantic.

538

539 Formation of the JMMC correlates with and can be explained by rejuvenation of pre-
540 existing structures of different ages. Oblique accommodation/deformation zones
541 including fracture zones defined the extent of the JMMC along the spreading axis. This
542 model provides a simple explanation for microplate-complex formation involving
543 control by both plate tectonic processes and structural inheritance.
544 Further work and data acquisition is required to fully understand the nature and
545 formation of the JMMC, Iceland and the Iceland-Faroe Ridge. All three components are
546 intrinsically interlinked and essential for understanding the tectonic and magmatic
547 evolution of the entire North Atlantic. Geophysical data are lacking especially in the
548 south of the JMMC, offshore northwest Iceland, and between Iceland and the Faroe
549 Islands. The most fundamental and perhaps economically important question is the
550 extent of continental crust underlying this region, a question that may require additional
551 marine surveys, re-interpretation of geochemical data and further drilling and sampling
552 in this area.

553

554 **ACKNOWLEDGEMENTS**

555 Christian Schiffer's postdoctoral fellowship at Durham University is funded by the
556 Carlsberg Foundation. Alexander Peace's postdoctoral fellowship at Memorial
557 University is funded by the Hibernia Project Geophysics Support Fund. We thank the
558 two anonymous reviewers for constructive comments that helped improving the paper.
559 We thank the North Atlantic research group for valuable inspiration during the 2016 and
560 2017 North Atlantic workshops at Durham University.

561

562 **REFERENCES**

- 563 Abdelmalak, M.M., Geoffroy, L., Angelier, J., Bonin, B., Callot, J.P., Gélard, J.P., and Aubourg, C.,
564 2012. Stress fields acting during lithosphere breakup above a melting mantle: A case
565 example in West Greenland, *Tectonophysics*, **581**, 132–143, doi:
566 10.1016/j.tecto.2011.11.020.
- 567 Abera, R., Wijk, J. van, and Axen, G., 2016. Formation of continental fragments: The Tamayo
568 Bank, Gulf of California, Mexico, *Geology*, **44**, 595–598, doi: 10.1130/G38123.1.
- 569 Abramovitz, T., and Thybo, H., 2000. Seismic images of Caledonian, lithosphere-scale collision
570 structures in the southeastern North Sea along Mona Lisa Profile 2, *Tectonophysics*,
571 **317**, 27–54, doi: 10.1016/S0040-1951(99)00266-8.
- 572 Andersen, T.B., Jamtveit, B., Dewey, J.F., and Swensson, E., 1991. Subduction and eduction of
573 continental crust: major mechanisms during continent-continent collision and orogenic
574 extensional collapse, a model based on the south Norwegian Caledonides, *Terra Nova*,
575 **3**, 303–310, doi: 10.1111/j.1365-3121.1991.tb00148.x.
- 576 Asimow, P.D., and Langmuir, C.H., 2003. The importance of water to oceanic mantle melting
577 regimes, *Nature*, **421**, 815–820, doi: 10.1038/nature01429.
- 578 Audet, P., and Bürgmann, R., 2011. Dominant role of tectonic inheritance in supercontinent
579 cycles, *Nat. Geosci.*, **4**, 184–187, doi: 10.1038/ngeo1080.
- 580 Balling, N., 2000. Deep seismic reflection evidence for ancient subduction and collision zones
581 within the continental lithosphere of northwestern Europe, *Tectonophysics*, **329**, 269–
582 300, doi: 10.1016/S0040-1951(00)00199-2.
- 583 Balling, N., 2013. The Lithosphere beneath Northern Europe: Structure and Evolution over
584 three billion years - Contributions from geophysical studies: Aarhus, Denmark.
- 585 Behn, M.D., and Lin, J., 2000. Segmentation in gravity and magnetic anomalies along the U.S.
586 East Coast passive margin: Implications for incipient structure of the oceanic
587 lithosphere, *J. Geophys. Res. Solid Earth*, **105**, 25769–25790, doi:
588 10.1029/2000JB900292.
- 589 Bellahsen, N., Leroy, S., Autin, J., Razin, P., d’Acremont, E., Sloan, H., Pik, R., Ahmed, A., and
590 Khanbari, K., 2013. Pre-existing oblique transfer zones and transfer/transform
591 relationships in continental margins: New insights from the southeastern Gulf of Aden,
592 Socotra Island, Yemen, *Tectonophysics*, **607**, 32–50, doi: 10.1016/j.tecto.2013.07.036.
- 593 Bergh, S., Corfu, F., Inge, P., Kullerud, K., Armitage, P.E.B., Zwaan, K.B., Erling, R., Holdsworth,
594 R., and Chattopadhy, A., 2012. Chapter 11: Was the Precambrian Basement of Western
595 Troms and Lofoten-Vesterålen in Northern Norway Linked to the Lewisian of Scotland?
596 A Comparison of Crustal Components, Tectonic Evolution and Amalgamation History,
597 in Sharkov, E. ed., *Tectonics - Recent Advances*, InTech.
- 598 Bingen, B., Andersson, J., Söderlund, U., and Möller, C., 2008. The Mesoproterozoic in the
599 Nordic countries, in *Episodes*, *Episodes*, p. 29–34.
- 600 Blischke, A., Gaina, C., Hopper, J.R., Péron-Pinvidic, G., Brandsdóttir, B., Guarnieri, P.,
601 Erlendsson, Ö., and Gunnarsson, K., 2016. The Jan Mayen microcontinent: an update
602 of its architecture, structural development and role during the transition from the Ægir
603 Ridge to the mid-oceanic Kolbeinsey Ridge, *NE Atl. Reg. Reappraisal Crustal Struct.*
604 *Tectonostratigraphy Magmat. Evol.*,.

- 605 Bott, M.H.P., 1988. A new look at the causes and consequences of the Icelandic hot-spot, *Geol.*
606 *Soc. Lond. Spec. Publ.*, **39**, 15–23, doi: 10.1144/GSL.SP.1988.039.01.03.
- 607 Bott, M.H.P., 1974. Deep Structure, Evolution and Origin of the Icelandic Transverse Ridge, *in*
608 Kristjansson, L. ed., *Geodynamics of Iceland and the North Atlantic Area*, NATO
609 Advanced Study Institutes Series 11, Springer Netherlands, p. 33–47.
- 610 Bott, M.H.P., 1983. The Crust Beneath the Iceland-Faeroe Ridge, *in* Bott, M.H.P., Saxov, S.,
611 Talwani, M., and Thiede, J. eds., *Structure and Development of the Greenland-Scotland*
612 *Ridge: New Methods and Concepts*, Springer US, Boston, MA, p. 63–75.
- 613 Bowling, J.C., and Harry, D.L., 2001. Geodynamic models of continental extension and the
614 formation of non-volcanic rifted continental margins, *Geol. Soc. Lond. Spec. Publ.*, **187**,
615 511–536, doi: 10.1144/GSL.SP.2001.187.01.25.
- 616 Brandsdóttir, B., Hooft, E.E.E., Mjelde, R., and Murai, Y., 2015. Origin and evolution of the
617 Kolbeinsey Ridge and Iceland Plateau, N-Atlantic, *Geochem. Geophys. Geosystems*, **16**,
618 612–634, doi: 10.1002/2014GC005540.
- 619 Breivik, A.J., Mjelde, R., Faleide, J.I., and Murai, Y., 2012. The eastern Jan Mayen
620 microcontinent volcanic margin, *Geophys. J. Int.*, **188**, 798–818, doi: 10.1111/j.1365-
621 246X.2011.05307.x.
- 622 Brown, E.L., and Leshner, C.E., 2014. North Atlantic magmatism controlled by temperature,
623 mantle composition and buoyancy, *Nat. Geosci.*, **7**, 820–824, doi: 10.1038/ngeo2264.
- 624 Buitter, S.J.H., and Torsvik, T.H., 2014. A review of Wilson Cycle plate margins: A role for mantle
625 plumes in continental break-up along sutures?, *Gondwana Res.*, **26**, 627–653, doi:
626 10.1016/j.gr.2014.02.007.
- 627 de Castro, D.L., Bezerra, F.H.R., Sousa, M.O.L., and Fuck, R.A., 2012. Influence of
628 Neoproterozoic tectonic fabric on the origin of the Potiguar Basin, northeastern Brazil
629 and its links with West Africa based on gravity and magnetic data, *J. Geodyn.*, **54**, 29–
630 42, doi: 10.1016/j.jog.2011.09.002.
- 631 Chalmers, J.A., and Pulvertaft, T.C.R., 2001. Development of the continental margins of the
632 Labrador Sea: a review, *Geol. Soc. Lond. Spec. Publ.*, **187**, 77–105, doi:
633 10.1144/GSL.SP.2001.187.01.05.
- 634 Chenin, P., Manatschal, G., Lavier, L.L., and Erratt, D., 2015. Assessing the impact of orogenic
635 inheritance on the architecture, timing and magmatic budget of the North Atlantic rift
636 system: a mapping approach, *J. Geol. Soc.*, **172**, 711–720, doi: 10.1144/jgs2014-139.
- 637 Christiansson, P., Faleide, J.I., and Berge, A.M., 2000. Crustal structure in the northern North
638 Sea: an integrated geophysical study, *Geol. Soc. Lond. Spec. Publ.*, **167**, 15–40, doi:
639 10.1144/GSL.SP.2000.167.01.02.
- 640 Collier, J.S., Sansom, V., Ishizuka, O., Taylor, R.N., Minshull, T.A., and Whitmarsh, R.B., 2008.
641 Age of Seychelles–India break-up, *Earth Planet. Sci. Lett.*, **272**, 264–277, doi:
642 10.1016/j.epsl.2008.04.045.
- 643 Darbyshire, F.A., White, R.S., and Priestley, K.F., 2000. Structure of the crust and uppermost
644 mantle of Iceland from a combined seismic and gravity study, *Earth Planet. Sci. Lett.*,
645 **181**, 409–428, doi: 10.1016/S0012-821X(00)00206-5.

- 646 Dasgupta, R., and Hirschmann, M.M., 2006. Melting in the Earth's deep upper mantle caused
647 by carbon dioxide, *Nature*, **440**, 659–662, doi: 10.1038/nature04612.
- 648 Dewey, J.F., 1988. Extensional collapse of orogens, *Tectonics*, **7**, 1123–1139, doi:
649 10.1029/TC007i006p01123.
- 650 Dewey, J.F., Ryan, P.D., and Andersen, T.B., 1993. Orogenic uplift and collapse, crustal
651 thickness, fabrics and metamorphic phase changes: the role of eclogites, *Geol. Soc.
652 Lond. Spec. Publ.*, **76**, 325–343, doi: 10.1144/GSL.SP.1993.076.01.16.
- 653 Dewey, J., and Spall, H., 1975. Pre-Mesozoic plate tectonics: How far back in Earth history can
654 the Wilson Cycle be extended?, *Geology*, **3**, 422–424, doi: 10.1130/0091-
655 7613(1975)3<422:PPTHFB>2.0.CO;2.
- 656 Doré, A.G., Lundin, E.R., Gibbons, A., Sømme, T.O., and Tørudbakken, B.O., 2015. Transform
657 margins of the Arctic: a synthesis and re-evaluation, *Geol. Soc. Lond. Spec. Publ.*, **431**,
658 SP431.8, doi: 10.1144/SP431.8.
- 659 Doré, A.G., Lundin, E.R., Jensen, L.N., Birkeland, Ø., Eliassen, P.E., and Fichler, C., 1999.
660 Principal tectonic events in the evolution of the northwest European Atlantic margin,
661 *Geol. Soc. Lond. Pet. Geol. Conf. Ser.*, **5**, 41–61, doi: 10.1144/0050041.
- 662 Doré, A.G., Lundin, E.R., Kuszniir, N.J., and Pascal, C., 2008. Potential mechanisms for the
663 genesis of Cenozoic domal structures on the NE Atlantic margin: pros, cons and some
664 new ideas, *Geol. Soc. Lond. Spec. Publ.*, **306**, 1–26, doi: 10.1144/SP306.1.
- 665 Dunlap, W.J., and Fossen, H., 1998. Early Paleozoic orogenic collapse, tectonic stability, and
666 late Paleozoic continental rifting revealed through thermochronology of K-feldspars,
667 southern Norway, *Tectonics*, **17**, 604–620, doi: 10.1029/98TC01603.
- 668 Duretz, T., Petri, B., Mohn, G., Schmalholz, S.M., Schenker, F.L., and Müntener, O., 2016. The
669 importance of structural softening for the evolution and architecture of passive
670 margins, *Sci. Rep.*, **6**, 38704, doi: 10.1038/srep38704.
- 671 Eldholm, O., and Grue, K., 1994. North Atlantic volcanic margins: Dimensions and production
672 rates, *J. Geophys. Res. Solid Earth*, **99**, 2955–2968, doi: 10.1029/93JB02879.
- 673 Ellis, D., and Stoker, M.S., 2014. The Faroe–Shetland Basin: a regional perspective from the
674 Paleocene to the present day and its relationship to the opening of the North Atlantic
675 Ocean, *Geol. Soc. Lond. Spec. Publ.*, **397**, SP397.1, doi: 10.1144/SP397.1.
- 676 Engen, Ø., Faleide, J.I., and Dyreng, T.K., 2008. Opening of the Fram Strait gateway: A review of
677 plate tectonic constraints, *Tectonophysics*, **450**, 51–69, doi:
678 10.1016/j.tecto.2008.01.002.
- 679 Fedorova, T., Jacoby, W.R., and Wallner, H., 2005. Crust–mantle transition and Moho model
680 for Iceland and surroundings from seismic, topography, and gravity data,
681 *Tectonophysics*, **396**, 119–140, doi: 10.1016/j.tecto.2004.11.004.
- 682 Fichler, C., Odinsen, T., Rueslåtten, H., Olesen, O., Vindstad, J.E., and Wienecke, S., 2011.
683 Crustal inhomogeneities in the Northern North Sea from potential field modeling:
684 Inherited structure and serpentinites?, *Tectonophysics*, **510**, 172–185, doi:
685 10.1016/j.tecto.2011.06.026.

- 686 Fichler, C., Rundhovde, E., Olesen, O., Sæther, B.M., Rueslåtten, H., Lundin, E., and Doré, A.G.,
687 1999. Regional tectonic interpretation of image enhanced gravity and magnetic data
688 covering the mid-Norwegian shelf and adjacent mainland, *Tectonophysics*, **306**, 183–
689 197, doi: 10.1016/S0040-1951(99)00057-8.
- 690 Fossen, H., 2010. Extensional tectonics in the North Atlantic Caledonides: a regional view, *Geol.*
691 *Soc. Lond. Spec. Publ.*, **335**, 767–793, doi: 10.1144/SP335.31.
- 692 Foulger, G.R., 2006. Older crust underlies Iceland, *Geophys. J. Int.*, **165**, 672–676, doi:
693 10.1111/j.1365-246X.2006.02941.x.
- 694 Foulger, G.R., and Anderson, D.L., 2005. A cool model for the Iceland hotspot, *J. Volcanol.*
695 *Geotherm. Res.*, **141**, 1–22, doi: 10.1016/j.jvolgeores.2004.10.007.
- 696 Foulger, G.R., Du, Z., and Julian, B.R., 2003. Icelandic-type crust, *Geophys. J. Int.*, **155**, 567–590,
697 doi: 10.1046/j.1365-246X.2003.02056.x.
- 698 Foulger, G.R., Natland, J.H., and Anderson, D.L., 2005a. A source for Icelandic magmas in
699 remelted Iapetus crust, *J. Volcanol. Geotherm. Res.*, **141**, 23–44, doi:
700 10.1016/j.jvolgeores.2004.10.006.
- 701 Foulger, G.R., Natland, J.H., Presnell, D.C., and Anderson, D.L., 2005b. Plates, Plumes, And
702 Paradigms: Boulder, Colorado.
- 703 Franke, D., 2013. Rifting, lithosphere breakup and volcanism: Comparison of magma-poor and
704 volcanic rifted margins, *Mar. Pet. Geol.*, **43**, 63–87, doi:
705 10.1016/j.marpetgeo.2012.11.003.
- 706 Frizon de Lamotte, D., Fourdan, B., Leleu, S., Leparmentier, F., and de Clarens, P., 2015. Style of
707 rifting and the stages of Pangea breakup, *Tectonics*, **34**, 2014TC003760, doi:
708 10.1002/2014TC003760.
- 709 Funck, T., Erlendsson, Ö., Geissler, W.H., Gradmann, S., Kimbell, G.S., McDermott, K., and
710 Petersen, U.K., 2016a. A review of the NE Atlantic conjugate margins based on seismic
711 refraction data, *Geol. Soc. Lond. Spec. Publ.*, **447**, SP447.9, doi: 10.1144/SP447.9.
- 712 Funck, T., Geissler, W.H., Kimbell, G.S., Gradmann, S., Erlendsson, Ö., McDermott, K., and
713 Petersen, U.K., 2016b. Moho and basement depth in the NE Atlantic Ocean based on
714 seismic refraction data and receiver functions, *Geol. Soc. Lond. Spec. Publ.*, **447**,
715 SP447.1, doi: 10.1144/SP447.1.
- 716 Gaina, C., Gernigon, L., and Ball, P., 2009. Palaeocene–Recent plate boundaries in the NE
717 Atlantic and the formation of the Jan Mayen microcontinent, *J. Geol. Soc.*, **166**, 601–
718 616, doi: 10.1144/0016-76492008-112.
- 719 Gaina, C., Müller, R.D., Brown, B.J., and Ishihara, T., 2003. Microcontinent formation around
720 Australia, *Geol. Soc. Am. Spec. Pap.*, **372**, 405–416, doi: 10.1130/0-8137-2372-8.405.
- 721 Garde, A.A., Hamilton, M.A., Chadwick, B., Grocott, J., and McCaffrey, K.J.W., 2002. The
722 Ketilidian orogen of South Greenland : geochronology, tectonics, magmatism, and
723 fore-arc accretion during Palaeoproterozoic oblique convergence, *Can. J. Earth Sci.*, **39**,
724 765–793, doi: 10.1139/E02-026.

- 725 Gee, D.G., Fossen, H., Henriksen, N., and Higgins, A.K., 2008. From the early Paleozoic
726 platforms of Baltica and Laurentia to the Caledonide orogen of Scandinavia and
727 Greenland, *in* Episodes, Episodes.
- 728 Geoffroy, L., Burov, E.B., and Werner, P., 2015. Volcanic passive margins: another way to break
729 up continents, *Sci. Rep.*, **5**, 14828, doi: 10.1038/srep14828.
- 730 Gernigon, L., Blischke, A., Nasuti, A., and Sand, M., 2015. Conjugate volcanic rifted margins,
731 sea-floor spreading and microcontinent: Insights from new high-resolution
732 aeromagnetic surveys in the Norway Basin, *Tectonics*, 2014TC003717, doi:
733 10.1002/2014TC003717.
- 734 Gernigon, L., Gaina, C., Olesen, O., Ball, P.J., Péron-Pinvidic, G., and Yamasaki, T., 2012. The
735 Norway Basin revisited: From continental breakup to spreading ridge extinction, *Mar.
736 Pet. Geol.*, **35**, 1–19, doi: 10.1016/j.marpetgeo.2012.02.015.
- 737 Gernigon, L., Lucazeau, F., Brigaud, F., Ringenbach, J.-C., Planke, S., and Le Gall, B., 2006. A
738 moderate melting model for the Vøring margin (Norway) based on structural
739 observations and a thermo-kinematical modelling: Implication for the meaning of the
740 lower crustal bodies, *Tectonophysics*, **412**, 255–278, doi: 10.1016/j.tecto.2005.10.038.
- 741 Gernigon, L., Ringenbach, J.C., Planke, S., and Le Gall, B., 2004. Deep structures and breakup
742 along volcanic rifted margins: Insights from integrated studies along the outer Vøring
743 Basin (Norway), *Mar. Pet. Geol.*, **21**, 363–372, doi: 10.1016/j.marpetgeo.2004.01.005.
- 744 Gerya, T., 2012. Origin and models of oceanic transform faults, *Tectonophysics*, **522–523**, 34–
745 54, doi: 10.1016/j.tecto.2011.07.006.
- 746 Gibson, G.M., Totterdell, J.M., White, L.T., Mitchell, C.H., Stacey, A.R., Morse, M.P., and
747 Whitaker, A., 2013. Pre-existing basement structure and its influence on continental
748 rifting and fracture zone development along Australia’s southern rifted margin, *J. Geol.
749 Soc.*, **170**, 365–377, doi: 10.1144/jgs2012-040.
- 750 Van Gool, J.A.M., Connelly, J.N., Marker, M., and Mengel, F.C., 2002. The Nagssugtoqidian
751 Orogen of West Greenland: tectonic evolution and regional correlations from a West
752 Greenland perspective, *Can. J. Earth Sci.*, **39**, 665–686, doi: 10.1139/e02-027.
- 753 Gorbatshev, R., and Bogdanova, S., 1993. Frontiers in the Baltic Shield, *Precambrian Res.*, **64**,
754 3–21, doi: 10.1016/0301-9268(93)90066-B.
- 755 Gorczyk, W., and Vogt, K., 2015. Tectonics and melting in intra-continental settings, *Gondwana
756 Res.*, **27**, 196–208, doi: 10.1016/j.gr.2013.09.021.
- 757 Greenhalgh, E.E., and Kusznir, N.J., 2007. Evidence for thin oceanic crust on the extinct Aegir
758 Ridge, Norwegian Basin, NE Atlantic derived from satellite gravity inversion, *Geophys.
759 Res. Lett.*, **34**, L06305, doi: 10.1029/2007GL029440.
- 760 Grocott, J., and McCaffrey, K., 2016. Basin Evolution and Destruction in an Early Proterozoic
761 Continental Margin: the Rinkian Fold-Thrust Belt of Central West Greenland, *J. Geol.
762 Soc.*,.
- 763 Gudlaugsson, S.T., Gunnarsson, K., Sand, M., and Skogseid, J., 1988. Tectonic and volcanic
764 events at the Jan Mayen Ridge microcontinent, *Geol. Soc. Lond. Spec. Publ.*, **39**, 85–93,
765 doi: 10.1144/GSL.SP.1988.039.01.09.

- 766 Hansen, J., Jerram, D.A., McCaffrey, K., and Passey, S.R., 2009. The onset of the North Atlantic
767 Igneous Province in a rifting perspective, *Geol. Mag.*, **146**, 309–325, doi:
768 10.1017/S0016756809006347.
- 769 Heron, P.J., Pysklywec, R.N., and Stephenson, R., 2016. Lasting mantle scars lead to perennial
770 plate tectonics, *Nat. Commun.*, **7**, 11834, doi: 10.1038/ncomms11834.
- 771 Hill, R.I., 1991. Starting plumes and continental break-up, *Earth Planet. Sci. Lett.*, **104**, 398–416,
772 doi: 10.1016/0012-821X(91)90218-7.
- 773 Hjartarson, Á., Erlendsson, Ö., and Blischke, A., 2017. The Greenland–Iceland–Faroe Ridge
774 Complex, *Geol. Soc. Lond. Spec. Publ.*, **447**, SP447.14, doi: 10.1144/SP447.14.
- 775 Holbrook, W.S., Larsen, H.C., Korenaga, J., Dahl-Jensen, T., Reid, I.D., Kelemen, P.B., Hopper,
776 J.R., Kent, G.M., Lizarralde, D., Bernstein, S., and Detrick, R.S., 2001. Mantle thermal
777 structure and active upwelling during continental breakup in the North Atlantic, *Earth
778 Planet. Sci. Lett.*, **190**, 251–266, doi: 10.1016/S0012-821X(01)00392-2.
- 779 Holdsworth, R.E., Butler, C.A., and Roberts, A.M., 1997. The recognition of reactivation during
780 continental deformation, *J. Geol. Soc.*, **154**, 73–78, doi: 10.1144/gsjgs.154.1.0073.
- 781 Hole, M.J., and Millett, J.M., 2016. Controls of Mantle Potential Temperature and Lithospheric
782 Thickness on Magmatism in the North Atlantic Igneous Province, *J. Petrol.*, **57**, 417–
783 436, doi: 10.1093/petrology/egw014.
- 784 Huerta, A., and Harry, D.L., 2012. Wilson cycles, tectonic inheritance, and rifting of the North
785 American Gulf of Mexico continental margin, *Geosphere*, **8**, 374, doi:
786 10.1130/GES00725.1.
- 787 Huismans, R., and Beaumont, C., 2011. Depth-dependent extension, two-stage breakup and
788 cratonic underplating at rifted margins., *Nature*, **473**, 74–78, doi:
789 10.1038/nature09988.
- 790 Indrevær, K., Bergh, S.G., Koehl, J.-B., Hansen, J.-A., Schermer, E.R., and Ingebrigtsen, A., 2013.
791 Post-Caledonian Brittle Fault Zones on the Hyperextended SW Barents Sea Margin:
792 New Insights into Onshore and Offshore Margin Architecture, *Nor. J. Geol.*, **93**.
- 793 Jamtveit, B., Brooker, R., Brooks, K., Larsen, L.M., and Pedersen, T., 2001. The water content of
794 olivines from the North Atlantic Volcanic Province, *Earth Planet. Sci. Lett.*, **186**, 401–
795 415, doi: 10.1016/S0012-821X(01)00256-4.
- 796 Jolivet, L., Raimbourg, H., Labrousse, L., Avigad, D., Leroy, Y., Austrheim, H., and Andersen,
797 T.B., 2005. Softening triggered by eclogitization, the first step toward exhumation
798 during continental subduction, *Earth Planet. Sci. Lett.*, **237**, 532–547, doi:
799 10.1016/j.epsl.2005.06.047.
- 800 Kandilarov, A., Mjelde, R., Pedersen, R.-B., Hellevang, B., Papenberg, C., Petersen, C.-J., Planert,
801 L., and Flueh, E., 2012. The northern boundary of the Jan Mayen microcontinent,
802 North Atlantic determined from ocean bottom seismic, multichannel seismic, and
803 gravity data, *Mar. Geophys. Res.*, **33**, 55–76, doi: 10.1007/s11001-012-9146-4.
- 804 Kerr, A., Ryan, B., Gower, C.F., Wardle, R.J., and Kerr, A., 1996. The Makkovik Province:
805 extension of the Ketilidian Mobile Belt in mainland North America, *Geol. Soc. Lond.
806 Spec. Publ.*, **112**, 155–177, doi: 10.1144/GSL.SP.1996.112.01.09.

- 807 King, S.D., and Anderson, D.L., 1998. Edge-driven convection, *Earth Planet. Sci. Lett.*, **160**, 289–
808 296, doi: 10.1016/S0012-821X(98)00089-2.
- 809 Kodaira, S., Mjelde, R., Gunnarsson, K., Shiobara, H., and Shimamura, H., 1997. Crustal
810 structure of the Kolbeinsey Ridge, North Atlantic, obtained by use of ocean bottom
811 seismographs, *J. Geophys. Res. Solid Earth*, **102**, 3131–3151, doi: 10.1029/96JB03487.
- 812 Kodaira, S., Mjelde, R., Gunnarsson, K., Shiobara, H., and Shimamura, H., 1998. Structure of the
813 Jan Mayen microcontinent and implications for its evolution, *Geophys. J. Int.*, **132**,
814 383–400.
- 815 Korenaga, J., 2004. Mantle mixing and continental breakup magmatism, *Earth Planet. Sci. Lett.*,
816 **218**, 463–473, doi: 10.1016/S0012-821X(03)00674-5.
- 817 Krabbendam, M., and Barr, T.D., 2000. Proterozoic orogens and the break-up of Gondwana:
818 why did some orogens not rift?, *J. Afr. Earth Sci.*, **31**, 35–49, doi: 10.1016/S0899-
819 5362(00)00071-3.
- 820 Kuvaas, B., and Kodaira, S., 1997. Research article: The formation of the Jan Mayen
821 microcontinent: the missing piece in the continental puzzle between the Møre-Vøring
822 Basins and East Greenland, *First Break*, **15**, 239–247, doi: 10.3997/1365-2397.1997008.
- 823 Larsen, L.M., Pedersen, A.K., Sorensen, E.V., Watt, W.S., and Duncan, R.A., 2013. Stratigraphy
824 and age of the Eocene Igertivâ Formation basalts, alkaline pebbles and sediments of
825 the Kap Dalton Group in the graben at Kap Dalton, East Greenland,.
- 826 Larsen, L.M., Pedersen, A.K., Tegner, C., and Duncan, R.A., 2014. Eocene to Miocene igneous
827 activity in NE Greenland: northward younging of magmatism along the East Greenland
828 margin, *J. Geol. Soc.*, **171**, 539–553, doi: 10.1144/jgs2013-118.
- 829 Larsen, H.C., and Saunders, A.D., 1998. Tectonism and volcanism at the southeast Greenland
830 rifted margin: a record of plume impact and later continental rapture, *Proc. Ocean
831 Drill. Program Sci. Results*, **152**, doi: 10.2973/odp.proc.sr.152.1998.
- 832 Lavier, L.L., and Manatschal, G., 2006. A mechanism to thin the continental lithosphere at
833 magma-poor margins., *Nature*, **440**, 324–328, doi: 10.1038/nature04608.
- 834 Leslie, A.G., Smith, M., and Soper, N.J., 2008. Laurentian margin evolution and the Caledonian
835 orogeny—A template for Scotland and East Greenland, *Geol. Soc. Am. Mem.*, **202**,
836 307–343, doi: 10.1130/2008.1202(13).
- 837 Lister, G.S., Etheridge, M.A., and Symonds, P.A., 1986. Detachment faulting and the evolution
838 of passive continental margins, *Geology*, **14**, 246, doi: 10.1130/0091-
839 7613(1986)14<246:DFATEO>2.0.CO;2.
- 840 Lorenz, H., Gee, D.G., Larionov, A.N., and Majka, J., 2012. The Grenville–Sveconorwegian
841 orogen in the high Arctic, *Geol. Mag.*, **149**, 875–891, doi:
842 10.1017/S0016756811001130.
- 843 Lundin, E.R., and Doré, A.G., 2011. Hyperextension, serpentization, and weakening: A new
844 paradigm for rifted margin compressional deformation, *Geology*, **39**, 347–350, doi:
845 10.1130/G31499.1.

- 846 Lundin, E.R., and Doré, A.G., 2005. NE Atlantic break-up: a re-examination of the Iceland
847 mantle plume model and the Atlantic–Arctic linkage, *Geol. Soc. Lond. Pet. Geol. Conf.
848 Ser.*, **6**, 739–754, doi: 10.1144/0060739.
- 849 Manatschal, G., Lavier, L., and Chenin, P., 2015. The role of inheritance in structuring
850 hyperextended rift systems: Some considerations based on observations and
851 numerical modeling, *Gondwana Res.*, **27**, 140–164, doi: 10.1016/j.gr.2014.08.006.
- 852 McBride, J.H., White, R.S., Smallwood, J.R., and England, R.W., 2004. Must magmatic intrusion
853 in the lower crust produce reflectivity?, *Tectonophysics*, **388**, 271–297, doi:
854 10.1016/j.tecto.2004.07.055.
- 855 Meyer, R., Wijk, J. van, and Gernigon, L., 2007. The North Atlantic Igneous Province: A review
856 of models for its formation, *Geol. Soc. Am. Spec. Pap.*, **430**, 525–552, doi:
857 10.1130/2007.2430(26).
- 858 Misra, A.A., 2016. *Tectonic Inheritance in Continental Rifts and Passive*: Springer.
- 859 Mittelstaedt, E., Ito, G., and Behn, M.D., 2008. Mid-ocean ridge jumps associated with hotspot
860 magmatism, *Earth Planet. Sci. Lett.*, **266**, 256–270, doi: 10.1016/j.epsl.2007.10.055.
- 861 Mjelde, R., Eckhoff, I., Solbakken, S., Kodaira, S., Shimamura, H., Gunnarsson, K., Nakanishi, A.,
862 and Shiobara, H., 2007a. Gravity and S-wave modelling across the Jan Mayen Ridge,
863 North Atlantic; implications for crustal lithology, *Mar. Geophys. Res.*, **28**, 27–41, doi:
864 10.1007/s11001-006-9012-3.
- 865 Mjelde, R., and Faleide, J.I., 2009. Variation of Icelandic and Hawaiian magmatism: evidence
866 for co-pulsation of mantle plumes?, *Mar. Geophys. Res.*, **30**, 61–72, doi:
867 10.1007/s11001-009-9066-0.
- 868 Mjelde, R., Goncharov, A., and Müller, R.D., 2013. The Moho: Boundary above upper mantle
869 peridotites or lower crustal eclogites? A global review and new interpretations for
870 passive margins, *Tectonophysics*, **609**, 636–650, doi: 10.1016/j.tecto.2012.03.001.
- 871 Mjelde, R., Kvarven, T., Faleide, J.I., and Thybo, H., 2016. Lower crustal high-velocity bodies
872 along North Atlantic passive margins, and their link to Caledonian suture zone
873 eclogites and Early Cenozoic magmatism, *Tectonophysics*, **670**, 16–29, doi:
874 10.1016/j.tecto.2015.11.021.
- 875 Mjelde, R., Raum, T., Murai, Y., and Takanami, T., 2007b. Continent–ocean-transitions: Review,
876 and a new tectono-magmatic model of the Vøring Plateau, NE Atlantic, *J. Geodyn.*, **43**,
877 374–392, doi: 10.1016/j.jog.2006.09.013.
- 878 Mohriak, W.U., and Rosendahl, B.R., 2003. Transform zones in the South Atlantic rifted
879 continental margins, *Geol. Soc. Lond. Spec. Publ.*, **210**, 211–228, doi:
880 10.1144/GSL.SP.2003.210.01.13.
- 881 Mosar, J., Lewis, G., and Torsvik, T., 2002. North Atlantic sea-floor spreading rates: implications
882 for the Tertiary development of inversion structures of the Norwegian–Greenland Sea,
883 *J. Geol. Soc.*, **159**, 503–515, doi: 10.1144/0016-764901-135.
- 884 Müller, R.D., Gaina, C., Roest, W.R., and Hansen, D.L., 2001. A recipe for microcontinent
885 formation, *Geology*, **29**, 203–206, doi: 10.1130/0091-
886 7613(2001)029<0203:ARFMF>2.0.CO;2.

- 887 Nemčok, M., Sinha, S.T., Doré, A.G., Lundin, E.R., Mascle, J., and Rybár, S., 2016. Mechanisms
888 of microcontinent release associated with wrenching-involved continental break-up; a
889 review, *Geol. Soc. Lond. Spec. Publ.*, **431**, SP431.14, doi: 10.1144/SP431.14.
- 890 Nichols, A.R.L., Carroll, M.R., and Höskuldsson, Á., 2002. Is the Iceland hot spot also wet?
891 Evidence from the water contents of undegassed submarine and subglacial pillow
892 basalts, *Earth Planet. Sci. Lett.*, **202**, 77–87, doi: 10.1016/S0012-821X(02)00758-6.
- 893 Nielsen, T.K., Larsen, H.C., and Hopper, J.R., 2002. Contrasting rifted margin styles south of
894 Greenland: implications for mantle plume dynamics, *Earth Planet. Sci. Lett.*, **200**, 271–
895 286, doi: 10.1016/S0012-821X(02)00616-7.
- 896 Nielsen, S.B., Stephenson, R., and Thomsen, E., 2007. Dynamics of Mid-Palaeocene North
897 Atlantic rifting linked with European intra-plate deformations, *Nature*, **450**, 1071–
898 1074, doi: 10.1038/nature06379.
- 899 Nirrengarten, M., Gernigon, L., and Manatschal, G., 2014. Lower crustal bodies in the Møre
900 volcanic rifted margin: Geophysical determination and geological implications,
901 *Tectonophysics*, **636**, 143–157, doi: 10.1016/j.tecto.2014.08.004.
- 902 Nunns, A., 1982. The Structure and Evolution of the Jan Mayen Ridge and Surrounding Regions:
903 Rifted Margins: Field Investigations of Margin Structure and Stratigraphy, **110**, 193–
904 208.
- 905 Oakey, G.N., and Chalmers, J. a, 2012. A new model for the Paleogene motion of Greenland
906 relative to North America : Plate reconstructions of the Davis Strait and Nares Strait
907 regions between Canada and Greenland, *J. Geophys. Res. Solid Earth*, **117**, 1–28, doi:
908 10.1029/2011JB008942.
- 909 Olafsson, I., Sundvor, E., Eldholm, O., and Grue, K., 1992. Møre Margin: Crustal structure from
910 analysis of Expanded Spread Profiles, *Mar. Geophys. Res.*, **14**, 137–162, doi:
911 10.1007/BF01204284.
- 912 Paquette, J., Sigmarsson, O., and Tiepolo, M., 2006. Continental basement under Iceland
913 revealed by old zircons, *AGU Fall Meet. Abstr.*, **33**.
- 914 Peace, A., McCaffrey, K., Imber, J., Phethean, J., Nowell, G., Gerdes, K., and Dempsey, E., 2016.
915 An evaluation of Mesozoic rift-related magmatism on the margins of the Labrador Sea:
916 Implications for rifting and passive margin asymmetry, *Geosphere*, **12**, 1701–1724, doi:
917 10.1130/GES01341.1.
- 918 Peron-Pinvidic, G., Gernigon, L., Gaina, C., and Ball, P., 2012. Insights from the Jan Mayen
919 system in the Norwegian–Greenland sea—I. Mapping of a microcontinent, *Geophys. J.*
920 *Int.*, **191**, 385–412, doi: 10.1111/j.1365-246X.2012.05639.x.
- 921 Peron-Pinvidic, G., and Manatschal, G., 2010. From microcontinents to extensional
922 allochthons: witnesses of how continents rift and break apart?, *Pet. Geosci.*, **16**, 189–
923 197, doi: 10.1144/1354-079309-903.
- 924 Peron-Pinvidic, G., Manatschal, G., and Osmundsen, P.T., 2013. Structural comparison of
925 archetypal Atlantic rifted margins: A review of observations and concepts, *Mar. Pet.*
926 *Geol.*, **43**, 21–47, doi: 10.1016/j.marpetgeo.2013.02.002.
- 927 Petersen, K.D., and Schiffer, C., 2016. Wilson cycle passive margins: Control of orogenic
928 inheritance on continental breakup, *Gondwana Res.*, doi: 10.1016/j.gr.2016.06.012.


- 929 Polat, A., Wang, L., and Appel, P.W.U., 2014. A review of structural patterns and melting
930 processes in the Archean craton of West Greenland: Evidence for crustal growth at
931 convergent plate margins as opposed to non-uniformitarian models, *Tectonophysics*,
932 **662**, 67–94, doi: 10.1016/j.tecto.2015.04.006.
- 933 Ren, S., Skogseid, J., and Eldholm, O., 1998. Late Cretaceous-Paleocene extension on the
934 Vøring Volcanic Margin, *Mar. Geophys. Res.*, **20**, 343–369, doi:
935 10.1023/A:1004554217069.
- 936 Rey, P., Vanderhaeghe, O., and Teyssier, C., 2001. Gravitational collapse of the continental
937 crust: definition, regimes and modes, *Tectonophysics*, **342**, 435–449, doi:
938 10.1016/S0040-1951(01)00174-3.
- 939 Reynisson, R.F., Ebbing, J., Lundin, E., and Osmundsen, P.T., 2010. Properties and distribution
940 of lower crustal bodies on the mid-Norwegian margin, *Geol. Soc. Lond. Pet. Geol. Conf.*
941 *Ser.*, **7**, 843–854, doi: 10.1144/0070843.
- 942 Richardson, K.R., Smallwood, J.R., White, R.S., Snyder, D.B., and Maguire, P.K.H., 1998. Crustal
943 structure beneath the Faroe Islands and the Faroe–Iceland Ridge, *Tectonophysics*, **300**,
944 159–180, doi: 10.1016/S0040-1951(98)00239-X.
- 945 Rickers, F., Fichtner, A., and Trampert, J., 2013. The Iceland–Jan Mayen plume system and its
946 impact on mantle dynamics in the North Atlantic region: Evidence from full-waveform
947 inversion, *Earth Planet. Sci. Lett.*, **367**, 39–51, doi: 10.1016/j.epsl.2013.02.022.
- 948 Roberts, D., 2003. The Scandinavian Caledonides: Event chronology, palaeogeographic settings
949 and likely modern analogues, *Tectonophysics*, **365**, 283–299, doi: 10.1016/S0040-
950 1951(03)00026-X.
- 951 Roest, W.R., and Srivastava, S.P., 1989. Sea-floor spreading in the Labrador Sea: a new
952 reconstruction, *Geology*, **17**, 1000–1003, doi: 10.1130/0091-
953 7613(1989)017<1000:SFSITL>2.3.CO;2.
- 954 Ryan, P.D., and Dewey, J.F., 1997. Continental eclogites and the Wilson Cycle, *J. Geol. Soc.*, **154**,
955 437–442, doi: 10.1144/gsjgs.154.3.0437.
- 956 Schiffer, C., Balling, N., Ebbing, J., Jacobsen, B.H., and Nielsen, S.B., 2016. Geophysical-
957 petrological modelling of the East Greenland Caledonides – Isostatic support from
958 crust and upper mantle, *Tectonophysics*,, doi: 10.1016/j.tecto.2016.06.023.
- 959 Schiffer, C., Balling, N., Jacobsen, B.H., Stephenson, R.A., and Nielsen, S.B., 2014. Seismological
960 evidence for a fossil subduction zone in the East Greenland Caledonides, *Geology*, **42**,
961 311–314, doi: 10.1130/G35244.1.
- 962 Schiffer, C., Jacobsen, B.H., Balling, N., Ebbing, J., and Nielsen, S.B., 2015a. The East Greenland
963 Caledonides—teleseismic signature, gravity and isostasy, *Geophys. J. Int.*, **203**, 1400–
964 1418, doi: 10.1093/gji/ggv373.
- 965 Schiffer, C., and Nielsen, S.B., 2016. Implications for anomalous mantle pressure and dynamic
966 topography from lithospheric stress patterns in the North Atlantic Realm, *J. Geodyn.*,
967 **98**, 53–69, doi: 10.1016/j.jog.2016.03.014.
- 968 Schiffer, C., Stephenson, R.A., Petersen, K.D., Nielsen, S.B., Jacobsen, B.H., Balling, N., and
969 Macdonald, D.I.M., 2015b. A sub-crustal piercing point for North Atlantic

- 970 reconstructions and tectonic implications, *Geology*, **43**, 1087–1090, doi:
971 10.1130/G37245.1.
- 972 Scrutton, R.A., 1976. Microcontinents and their Significance, in Lake, C. ed., *Geodynamics:*
973 *Progress and Prospects*, American Geophysical Union, p. 177–189.
- 974 Seton, M., Müller, R.D., Zahirovic, S., Gaina, C., Torsvik, T., Shephard, G., Talsma, A., Gurnis, M.,
975 Turner, M., Maus, S., and Chandler, M., 2012. Global continental and ocean basin
976 reconstructions since 200 Ma, *Earth-Sci. Rev.*, **113**, 212–270, doi:
977 10.1016/j.earscirev.2012.03.002.
- 978 Simon, K., Huismans, R.S., and Beaumont, C., 2009. Dynamical modelling of lithospheric
979 extension and small-scale convection: Implications for magmatism during the
980 formation of volcanic rifted margins, *Geophys. J. Int.*, **176**, 327–350, doi:
981 10.1111/j.1365-246X.2008.03891.x.
- 982 Sinha, S.T., Nemčok, M., Choudhuri, M., Sinha, N., and Rao, D.P., 2016. The role of break-up
983 localization in microcontinent separation along a strike-slip margin: the East India–Elan
984 Bank case study, *Geol. Soc. Lond. Spec. Publ.*, **431**, 95–123, doi: 10.1144/SP431.5.
- 985 Skogseid, J., Planke, S., Faleide, J.I., Pedersen, T., Eldholm, O., and Neverdal, F., 2000. NE
986 Atlantic continental rifting and volcanic margin formation, *Geol. Soc. Lond. Spec. Publ.*,
987 **167**, 295–326, doi: 10.1144/GSL.SP.2000.167.01.12.
- 988 Slagstad, T., Roberts, N.M.W., and Kulakov, E., 2017. Linking orogenesis across a
989 supercontinent; the Grenvillian and Sveconorwegian margins on Rodinia, *Gondwana*
990 *Res.*, **44**, 109–115, doi: 10.1016/j.gr.2016.12.007.
- 991 Smallwood, J.R., Staples, R.K., Richardson, K.R., White, R.S., Brandsdóttir, B., Einarsson, P.,
992 England, R., Hobbs, R., Maguire, P., McBride, J., Menke, W., Minshull, T., Snyder, D.,
993 and Worthington, M., 1999. Crust generated above the Iceland mantle plume: From
994 continental rift to oceanic spreading center, *J. Geophys. Res. Solid Earth*, **104**, 22885–
995 22902.
- 996 Smallwood, J.R., and White, R.S., 2002. Ridge-plume interaction in the North Atlantic and its
997 influence on continental breakup and seafloor spreading, *Geol. Soc. Lond. Spec. Publ.*,
998 **197**, 15–37, doi: 10.1144/GSL.SP.2002.197.01.02.
- 999 Snyder, D.B., and Flack, C.A., 1990. A Caledonian age for reflectors within the mantle
1000 lithosphere north and west of Scotland, *Tectonics*, **9**, 903–922, doi:
1001 10.1029/TC009i004p00903.
- 1002 Srivastava, S.P., 1978. Evolution of the Labrador Sea and its bearing on the early evolution of
1003 the North Atlantic, *Geophys. J. Int.*, **52**, 313–357, doi: 10.1111/j.1365-
1004 246X.1978.tb04235.x.
- 1005 Staples, R.K., White, R.S., Brandsdóttir, B., Menke, W., Maguire, P.K.H., and McBride, J.H.,
1006 1997. Färoe-Iceland Ridge Experiment 1. Crustal structure of northeastern Iceland, *J.*
1007 *Geophys. Res. Solid Earth*, **102**, 7849–7866, doi: 10.1029/96JB03911.
- 1008 St-Onge, M.R., Gool, J.A.M. Van, Garde, A.A., and Scott, D.J., 2009. Correlation of Archaean and
1009 Palaeoproterozoic units between northeastern Canada and western Greenland:
1010 constraining the pre-collisional upper plate accretionary history of the Trans-Hudson
1011 orogen, *Geol. Soc. Lond. Spec. Publ.*, **318**, 193–235, doi: 10.1144/SP318.7.

- 1012 Sutherland, R., Davey, F., and Beavan, J., 2000. Plate boundary deformation in South Island,
1013 New Zealand, is related to inherited lithospheric structure, *Earth Planet. Sci. Lett.*, **177**,
1014 141–151, doi: 10.1016/S0012-821X(00)00043-1.
- 1015 Svartman Dias, A.E., Lavier, L.L., and Hayman, N.W., 2015. Conjugate rifted margins width and
1016 asymmetry: The interplay between lithospheric strength and thermomechanical
1017 processes, *J. Geophys. Res. Solid Earth*, **120**, 2015JB012074, doi:
1018 10.1002/2015JB012074.
- 1019 Talwani, M., and Eldholm, O., 1977. Evolution of the Norwegian-Greenland Sea, *Bull. Geol. Soc.*
1020 *Am.*, **88**, 969–999, doi: 10.1130/0016-7606(1977)88<969:EOTNS>2.0.CO;2.
- 1021 Tate, M.P., 1992. The Clare Lineament: a relic transform fault west of Ireland, *Geol. Soc. Lond.*
1022 *Spec. Publ.*, **62**, 375–384, doi: 10.1144/GSL.SP.1992.062.01.28.
- 1023 Taylor, B., Goodliffe, A., and Martinez, F., 2009. Initiation of transform faults at rifted
1024 continental margins, *Comptes Rendus Geosci.*, **341**, 428–438, doi:
1025 10.1016/j.crte.2008.08.010.
- 1026 Tegner, C., Brooks, C.K., Duncan, R.A., Heister, L.E., and Bernstein, S., 2008. 40Ar–39Ar ages of
1027 intrusions in East Greenland: Rift-to-drift transition over the Iceland hotspot, *Lithos*,
1028 **101**, 480–500, doi: 10.1016/j.lithos.2007.09.001.
- 1029 Tetreault, J.L., and Buitter, S.J.H., 2014. Future accreted terranes: a compilation of island arcs,
1030 oceanic plateaus, submarine ridges, seamounts, and continental fragments, *Solid*
1031 *Earth*, **5**, 1243–1275, doi: 10.5194/se-5-1243-2014.
- 1032 Thomas, W.A., 2006. Tectonic inheritance at a continental margin, *GSA Today*, **16**, 4–11.
- 1033 Thybo, H., and Artemieva, I.M., 2013. Moho and magmatic underplating in continental
1034 lithosphere, *Tectonophysics*, **609**, 605–619, doi: 10.1016/j.tecto.2013.05.032.
- 1035 Tommasi, A., Knoll, M., Vauchez, A., Signorelli, J.W., Thoraval, C., and Logé, R., 2009. Structural
1036 reactivation in plate tectonics controlled by olivine crystal anisotropy, *Nat. Geosci.*, **2**,
1037 423–427, doi: 10.1038/ngeo528.
- 1038 Tommasi, A., and Vauchez, A., 2015. Heterogeneity and anisotropy in the lithospheric mantle,
1039 *Tectonophysics*, **661**, 11–37, doi: 10.1016/j.tecto.2015.07.026.
- 1040 Torsvik, T.H., Amundsen, H.E.F., Trønnes, R.G., Doubrovine, P. V., Gaina, C., Kuznir, N.J.,
1041 Steinberger, B., Corfu, F., Ashwal, L.D., Griffin, W.L., Werner, S.C., and Jamtveit, B.,
1042 2015. Continental crust beneath southeast Iceland, *Proc. Natl. Acad. Sci.*, 201423099,
1043 doi: 10.1073/pnas.1423099112.
- 1044 Torsvik, T.H., Carlos, D., Mosar, J., Cocks, L.R.M., and Malme, T., 2002. Global reconstructions
1045 and North Atlantic palaeogeography 400 Ma to Recent., in *BATLAS – Mid Norway plate*
1046 *reconstructions atlas with global and Atlantic perspectives.*, Geological Survey of
1047 Norway, p. 18–39.
- 1048 Upton, B.G.J., 1988. History of Tertiary igneous activity in the N Atlantic borderlands, *Geol. Soc.*
1049 *Lond. Spec. Publ.*, **39**, 429–453, doi: 10.1144/GSL.SP.1988.039.01.38.
- 1050 Vauchez, A., Barruol, G., and Tommasi, A., 1997. Why do continents break-up parallel to
1051 ancient orogenic belts?, *Terra Nova*, **9**, 62–66, doi: 10.1111/j.1365-
1052 3121.1997.tb00003.x.

- 1053 Vauchez, A., Tommasi, A., and Barruol, G., 1998. Rheological heterogeneity, mechanical
1054 anisotropy and deformation of the continental lithosphere, *Tectonophysics*, **296**, 61–
1055 86, doi: 10.1016/S0040-1951(98)00137-1.
- 1056 van der Velden, A.J., and Cook, F.A., 2005. Relict subduction zones in Canada, *J. Geophys. Res.*
1057 *Solid Earth*, **110**, n/a–n/a, doi: 10.1029/2004JB003333.
- 1058 Vink, G.E., 1984. A hotspot model for Iceland and the Vøring Plateau, *J. Geophys. Res. Solid*
1059 *Earth*, **89**, 9949–9959, doi: 10.1029/JB089iB12p09949.
- 1060 Vogt, P.R., Ostenso, N.A., and Johnson, G.L., 1970. Magnetic and bathymetric data bearing on
1061 sea-floor spreading north of Iceland, *J. Geophys. Res.*, **75**, 903–920, doi:
1062 10.1029/JB075i005p00903.
- 1063 Vogt, P.R., Perry, R.K., Feden, R.H., Fleming, H.S., and Cherkis, N.Z., 1981. The Greenland—
1064 Norwegian Sea and Iceland Environment: Geology and Geophysics, in Nairn, A.E.M., Jr,
1065 M.C., and Stehli, F.G. eds., *The Arctic Ocean*, Springer US, p. 493–598.
- 1066 Wangen, M., Mjelde, R., and Faleide, J.I., 2011. The extension of the Vøring margin (NE
1067 Atlantic) in case of different degrees of magmatic underplating, *Basin Res.*, **23**, 83–100,
1068 doi: 10.1111/j.1365-2117.2010.00467.x.
- 1069 Wardle, R.J., James, D.T., Scott, D.J., and Hall, J., 2002. The southeastern Churchill Province:
1070 synthesis of a Paleoproterozoic transpressional orogen, *Can. J. Earth Sci.*, **39**, 639–663,
1071 doi: 10.1139/e02-004.
- 1072 Warner, M., Morgan, J., Barton, P., Morgan, P., Price, C., and Jones, K., 1996. Seismic
1073 reflections from the mantle represent relict subduction zones within the continental
1074 lithosphere, *Geology*, **24**, 39–42, doi: 10.1130/0091-
1075 7613(1996)024<0039:SRFTMR>2.3.CO;2.
- 1076 Whalen, L., Gazel, E., Vidito, C., Puffer, J., Bizimis, M., Henika, W., and Caddick, M.J., 2015.
1077 Supercontinental inheritance and its influence on supercontinental breakup: The
1078 Central Atlantic Magmatic Province and the breakup of Pangea, *Geochem. Geophys.*
1079 *Geosystems*, **16**, 3532–3554, doi: 10.1002/2015GC005885.
- 1080 White, R., and McKenzie, D., 1989. Magmatism at rift zones: The generation of volcanic
1081 continental margins and flood basalts, *J. Geophys. Res.*, **94**, 7685, doi:
1082 10.1029/JB094iB06p07685.
- 1083 White, R.S., Smith, L.K., Roberts, a W., Christie, P. a F., Kuszniir, N.J., Roberts, a M., Healy, D.,
1084 Spitzer, R., Chappell, a, Eccles, J.D., Fletcher, R., Hurst, N., Lunnon, Z., Parkin, C.J., et
1085 al., 2008. Lower-crustal intrusion on the North Atlantic continental margin., *Nature*,
1086 **452**, 460–464, doi: 10.1038/nature06687.
- 1087 Whittaker, J.M., Williams, S.E., Halpin, J.A., Wild, T.J., Stilwell, J.D., Jourdan, F., and Daczko,
1088 N.R., 2016. Eastern Indian Ocean microcontinent formation driven by plate motion
1089 changes, *Earth Planet. Sci. Lett.*, **454**, 203–212, doi: 10.1016/j.epsl.2016.09.019.
- 1090 van Wijk, J.W., Huismans, R.S., ter Voorde, M., and Cloetingh, S. a. P.L., 2001. Melt generation
1091 at volcanic continental margins: No need for a mantle plume?, *Geophys. Res. Lett.*, **28**,
1092 3995–3998, doi: 10.1029/2000GL012848.
- 1093 Wilson, J.T., 1966. Did the Atlantic Close and then Re-Open?, *Nature*, **211**, 676–681, doi:
1094 10.1038/211676a0.

- 1095 Yamasaki, T., and Gernigon, L., 2010. Redistribution of the lithosphere deformation by the
1096 emplacement of underplated mafic bodies: implications for microcontinent formation,
1097 *J. Geol. Soc.*, **167**, 961–971, doi: 10.1144/0016-76492010-027.
- 1098 Zhang, J., and Green, H.W., 2007. Experimental Investigation of Eclogite Rheology and Its
1099 Fabrics at High Temperature and Pressure, *J. Metamorph. Geol.*, **25**, 97–115, doi:
1100 10.1111/j.1525-1314.2006.00684.x.
- 1101 Ziegler, P.A. (Ed.), 1990. Geological Atlas of Western and Central Europe: Geological Society of
1102 London, The Hague.
- 1103




1105 **Figure 1**

1106 Bathymetric map of the present-day North Atlantic. Bathymetry from the General
 1107 Bathymetric Chart of the Oceans (GEBCO). Major oceanic fracture zones after Dore *et*
 1108 *al.* (2008), Mid Ocean Ridges from Seton *et al.* (2012), microcontinents from Torsvik *et*
 1109 *al.* (2015). Greenland-Iceland-Faroe Ridge (GIFR) consists of the Greenland-Iceland
 1110 Ridge, the Iceland Plateau and the Iceland-Faroe Ridge. The position of the Iceland
 1111 Faroe Fracture Zone is stippled, but its existence and nature is debated (see text). AO =
 1112 Arctic Ocean; AR = Aegir Ridge; BB = Baffin Bay; BFZ = Bight Fracture Zone; BI =
 1113 Baffin Island; BR = Britain; BS = Barents Sea; CGFZ = Charlie-Gibbs Fracture Zone;
 1114 DS = Davis Strait; EB = Eurasia basin; EI = Ellesmere Island; EJMFBZ = East Jan
 1115 Mayen Fracture Zone; GIR = Greenland-Iceland Ridge; GR = Greenland; IC – Iceland;
 1116 IFFZ = Iceland-Faroe Fracture Zone; IFR = Iceland-Faroe Ridge; IR = Ireland; KR =
 1117 Kolbeinsey Ridge; LA = Labrador; LS = Labrador Sea; NF = Newfoundland; NS =
 1118 Nares Strait; RP = Rockall Plateau; RR = Reykjanes Ridge; SC = Scandinavia; SFZ =

1119 Senja Fracture Zone: SF = Svecofennian; SI = Shetland Islands; SV = Svalbard;
 1120 WJMFZ = West Jan Mayen Fracture Zone.

1121

1122



1123


1124 Figure 2

1125 Overview map of the present-day North Atlantic. Seafloor age from Seton *et al.* (2012),
 1126 major oceanic fracture zones after Doré *et al.* (2008), distribution of igneous rocks of
 1127 the North Atlantic Igneous Province after Upton (1988), Larsen & Saunders (1998),
 1128 Abdelmalak *et al.* (2012), Precambrian basement terranes after Balling (2000) and
 1129 Indrevær *et al.* (2013) – Scandinavia, St-Onge *et al.* (2009) – Greenland and
 1130 northeastern Canada. Caledonian Deformation Front after Skogseid *et al.* (2000) and
 1131 Gee *et al.* (2008). K = Karelian; KE = Ketilian Orogen; LK = Lapland-Kola; NAC =

1132 North Atlantic Craton; NO = Nagssugtoqidian Orogen; RO = Rinkian Orogen; SF =
1133 Svecofennian; SN = Sveconorwegian; TIB = Transscandinavian Igneous Belt.

1134

1135



1136


1137 Figure 3

1138 Bathymetry (a), free air gravity (b) and magnetic anomaly (c) maps of the Norway
1139 Basin, the Jan Mayen microplate complex (JMMC), Iceland, the Iceland-Faroe Ridge
1140 and surrounding conjugate margins (modified after Gernigon *et al.* 2015). The
1141 bathymetric map illustrates the special physiological nature of the JMMC, coinciding

1142 with large free air gravity anomalies. Magnetic anomalies within the boundaries of the
 1143 JMMC are weak. This is in large contrast to the adjacent Norway Basin, which shows
 1144 clear magnetic spreading anomalies, and gravity and topographic anomalies that
 1145 evidence the “fan-shaped” spreading along the extinct Aegir Ridge. There are vague
 1146 indications in bathymetry, gravity and magnetic data for the existence of a lineament
 1147 stretching from the south of the JMMC to the Faroe-Shetland Basin, possibly the IFFZ
 1148 (Blischke *et al.*, 2016), but the data does not provide indisputable evidence for the
 1149 existence and the nature of such.

1150

1151



1152


1153 Figure 4

1154 Model for the formation of the Batavia and Gulden Draak microcontinents in the Indian
 1155 Ocean proposed by Whittaker *et al.* (2016). Initial seafloor spreading occurred
 1156 perpendicular to the regional plate motions, including the Wallaby-Zenith Fracture Zone
 1157 (WZFZ). A reconfiguration of plate motions oblique to the developed spreading axes
 1158 locked the fracture zone, which forced the southern spreading axis to relocate onto a
 1159 new axis. The new spreading isolates continental fragments (microcontinents) and
 1160 seafloor spreading separates these from the Indian plate. Large arrows indicate plate
 1161 motions. Arrows along spreading ridges indicate the spreading direction. Dots with
 1162 arrows indicate the transpressional regime along the former fracture zone.

1163

1164

1165




1166

1167 Figure 5

1168 Application of the model of Whittaker *et al.* (2016) to the formation of the Jan Mayen
 1169 microplate complex. The original model was developed to explain microcontinent
 1170 separation between Greater India and Australia. (a) NW-SE plate motion between
 1171 Greenland and Europe with the Iceland-Faroe accommodation zone (IFAZ) as a diffuse
 1172 zone accommodating relative motion between the Reykjanes ridge (RR) and Aegir ridge
 1173 (AR). Continental rifting and extension occurs along the lithospheric weakness (East
 1174 Greenland fossil subduction zone) (b) Plate tectonic reorganisations result in W-E
 1175 motion between Greenland and Europe locking up the Iceland-Faroe accommodation
 1176 zone. The Reykjanes ridge diverts towards the north following the lithospheric
 1177 weakness. (c) Seafloor spreading develops along the Kolbeinsey ridge (KR) breaking
 1178 the Jan Mayen Microplate off from Greenland. The JMMC rotates counterclockwise.
 1179 Seafloor spreading on the Aegir ridge is abandoned.

1180



1182

1183 Figure 6:

1184 Separation of the Jan Mayen microplate complex from Greenland. Palaeogeographic
 1185 reconstructions from Seton *et al.* (2012). 100 Ma: The Caledonian Orogen experienced
 1186 extensional collapse and multiple rift phases. Fossil subduction zones are still preserved,
 1187 though possibly deformed. 50 Ma: Seafloor spreading in the North Atlantic separates
 1188 Greenland from Europe with NW-SE plate motions. Breakup in the NE Atlantic occurs
 1189 along the Iapetus suture, which deforms. 40 Ma: Plate motions change from NW-SE to
 1190 W-E, which causes transpression on the Iceland-Faroe accommodation zone. The
 1191 Reykjanes ridge spreading centre develops towards the north, following lithospheric

1192 weaknesses along the East Greenland fossil subduction zone. 20 Ma: The newly formed
1193 Kolbeinsey ridge is almost entirely developed, separating the Jan Mayen Microplate
1194 Complex from Greenland. The fossil subduction zone in Central East Greenland is
1195 highly deformed, whereas it is mainly preserved further north. The Aegir Ridge is
1196 successively abandoned. 0 Ma: Fossil subduction zones are still preserved in East
1197 Greenland, northern Scotland and the Danish North Sea sector (Central Fjord structure -
1198 CF, Flannan reflector - FL, Mona Lisa structure - ML). In Norway and south-central
1199 East Greenland the fossil subduction zone has been destroyed and deformed. It now
1200 forms high-seismic-velocity lower crustal bodies that are possible eclogite HVLCBs
1201 mapped in magenta and orange).

1202